

UB-ECM-PF-97/28
hep-th/9712125
to appear in Nucl. Phys. B

Cohomological analysis of bosonic D-strings and 2d sigma models coupled to abelian gauge fields

Friedemann Brandt, Joaquim Gomis, Joan Simón

Departament ECM, Facultat de Física, Universitat de Barcelona and Institut de Física d'Altes Energies, Diagonal 647, E-08028 Barcelona, Spain.

E-mail: brandt@itp.uni-hannover.de, gomis@ecm.ub.es, jsimon@ecm.ub.es.

Abstract

We analyse completely the BRST cohomology on local functionals for two dimensional sigma models coupled to abelian world sheet gauge fields, including effective bosonic D-string models described by Born-Infeld actions. In particular we prove that the rigid symmetries of such models are exhausted by the solutions to generalized Killing vector equations which we have presented recently, and provide all the consistent first order deformations and candidate gauge anomalies of the models under study. For appropriate target space geometries we find nontrivial deformations both of the abelian gauge transformations and of the world sheet diffeomorphisms, and antifield dependent candidate anomalies for both types of symmetries separately, as well as mixed ones.

PACS numbers: 02.90.+p, 11.10.Kk, 11.25.-w, 11.30.-j

Keywords: D-strings, BRST cohomology, rigid symmetries, conservation laws, consistent deformations, anomalies

1 Introduction

D-branes [1] play a crucial role in recent advances towards understanding nonperturbative properties of string and supersymmetric quantum field theories. The low energy effective action of a D-brane involves a Born–Infeld type action containing the coordinates of the brane and a $U(1)$ world-volume gauge field. In the case of IIB D-branes, it has been recently suggested that, for a manifestly $SL(2, Z)$ invariant formulation, the action should contain a world-volume field for every background gauge potential [2, 3]. In the case of the D-string one then should introduce two $U(1)$ gauge fields on the world-sheet.

Inspired by these developments, we consider in this paper a class of models including (effective) bosonic D-strings described by Born–Infeld type actions, but not restricted to them. Rather, the models under study are characterized only by their field content, gauge symmetries and a locality requirement. Specifically, we consider models described in terms of a set of world-sheet scalar fields X^M (“target space coordinates”), a set of world-sheet gauge fields A_μ^a , and an auxiliary world-sheet metric $\gamma_{\mu\nu}$. The action functionals for such models are required to be local, i.e. polynomial in derivatives of the fields, and invariant under world-sheet diffeomorphisms, local Weyl transformations of $\gamma_{\mu\nu}$ and abelian gauge transformations of the A_μ^a (the local Weyl symmetry is spurious in the sense that it can be avoided if one works with two “Beltrami variables” h_{++} , h_{--} instead of the three components of $\gamma_{\mu\nu}$, see section 3). The classification of all action functionals satisfying these requirements will be part of our results. This includes Born–Infeld type actions, as they can be written in a form which is local in the above sense by means of scalar auxiliary fields counting among the X^M , see section 4.

The main purpose of the paper is to investigate these models with regard to their symmetries. Specifically, we aim at a classification of their rigid (= global) symmetries and dynamical conservation laws, and of the possible anomalies and nontrivial continuous deformations of their gauge (= local) symmetries. Studying such deformations covers the determination of possible gauge invariant interactions between world sheet gauge fields and the target fields X^M , and has useful applications within the renormalization program for effective gauge theories [4].

These issues are analysed systematically by computing the BRST cohomology in the space of local functionals of the fields and their antifields. The rigid symmetries and dynamical conservation laws arise from this cohomology at negative ghost numbers [5], the action functionals and consistent (first order) deformations at ghost number 0 [6], and the candidate anomalies at ghost number 1 [7]. In fact we will compute the cohomology at all ghost numbers and discuss afterwards the results in detail for ghost numbers ≤ 1 .

We have organized the paper as follows. In section 2 we describe the models and the cohomological problem to be studied. In section 3 we perform the first part of the cohomological analysis for which the explicit knowledge of a classical action is not needed. The action functionals themselves are then determined in section 4. We then complete the cohomological analysis and summarize the result in a compact form for

all ghost numbers in section 5 (the computation itself is relegated to an appendix). In section 6 we spell out explicitly the physically most important solutions (those with ghost numbers ≤ 1) and comment on them. To illustrate these results we discuss in section 7 a class of models with only one abelian gauge field. The paper is ended by some conclusions in section 8, and three technical appendices.

2 The cohomological problem

As mentioned already, the field content of the models we are going to study is given by the world-sheet metric $\gamma_{\mu\nu}$ ($\mu = +, -$), a set of scalar fields X^M – some of which may be auxiliary fields – and a set of abelian gauge fields A_μ^a . The number of scalar and gauge fields is not specified, i.e. our approach covers models with any number of such fields. We impose invariance under world-sheet diffeomorphisms, local Weyl transformations of the world-sheet metric, and abelian gauge transformations of the A_μ^a . The ghost fields associated with these gauge symmetries are denoted by ξ^μ (diffeomorphism ghosts), c (Weyl ghost) and C^a (abelian ghosts) respectively. This fixes the field content to

$$\{\Phi^A\} = \{\gamma_{\mu\nu}, X^M, A_\mu^a, \xi^\mu, c, C^a\}.$$

The BRST transformations of these fields corresponding to the above mentioned gauge symmetries are

$$\begin{aligned} s\gamma_{\mu\nu} &= \xi^\rho \partial_\rho \gamma_{\mu\nu} + \partial_\mu \xi^\rho \cdot \gamma_{\rho\nu} + \partial_\nu \xi^\rho \cdot \gamma_{\mu\rho} + c \gamma_{\mu\nu} \\ sX^M &= \xi^\mu \partial_\mu X^M \\ sA_\mu^a &= \xi^\nu \partial_\nu A_\mu^a + \partial_\mu \xi^\nu \cdot A_\nu^a + \partial_\mu C^a \\ s\xi^\mu &= \xi^\nu \partial_\nu \xi^\mu \\ sc &= \xi^\mu \partial_\mu c \\ sC^a &= \xi^\mu \partial_\mu C^a. \end{aligned} \tag{2.1}$$

According to the standard prescription of the field-antifield formalism [8, 9, 10], the BRST transformations of the antifields are given by

$$s\Phi_A^* \equiv (\mathcal{S}, \Phi_A^*) = \frac{\delta^R \mathcal{S}}{\delta \Phi^A} \tag{2.2}$$

where $\mathcal{S}$ is the proper solution of the (classical) master equation. As the gauge algebra is closed, $\mathcal{S}$ reads in our case simply

$$\mathcal{S} = \int d^2\sigma \left[L_0 - (s\Phi^A) \Phi_A^* \right] \tag{2.3}$$

where L_0 is the integrand of a classical action with (only) the above gauge symmetries. For later purposes we decompose the BRST-transformations into parts with different so-called antighost numbers. In this case the decomposition has only two pieces,

denoted by δ and γ respectively where δ is the field theoretical Koszul–Tate differential [11] which acts nontrivially only on the antifields,

$$s = \delta + \gamma, \quad s\Phi^A = \gamma\Phi^A, \quad s\Phi_A^* = (\delta + \gamma)\Phi_A^*. \quad (2.4)$$

Only δ depends on the explicit form of L_0 ,

$$\begin{aligned} \delta\phi_i^* &= \frac{\hat{\partial}L_0}{\hat{\partial}\phi^i} \quad \text{for} \quad \{\phi^i\} = \{X^M, A_\mu^a, \gamma_{\mu\nu}\} \\ \delta\xi_\mu^* &= X_M^* \partial_\mu X^M + \gamma^{\rho\sigma*} \partial_\mu \gamma_{\rho\sigma} + A_a^{\nu*} \partial_\mu A_\nu^a + \partial_\nu (2\gamma_{\mu\rho} \gamma^{\nu\rho*} + A_\mu^a A_a^{\nu*}) \\ \delta c^* &= \gamma_{\mu\nu} \gamma^{\mu\nu*} \\ \delta C_a^* &= -\partial_\mu A_a^{\mu*} \end{aligned} \quad (2.5)$$

where $\hat{\partial}L_0/\hat{\partial}\phi^i$ is the Euler–Lagrange derivative of L_0 with respect to ϕ^i . In contrast, γ is determined already by the gauge symmetries only,

$$\begin{aligned} \gamma X_M^* &= \partial_\mu (\xi^\mu X_M^*) \\ \gamma A_a^{\mu*} &= \partial_\nu (\xi^\nu A_a^{\mu*}) - \partial_\nu \xi^\mu \cdot A_a^{\nu*} \\ \gamma \gamma^{\mu\nu*} &= \partial_\rho (\xi^\rho \gamma^{\mu\nu*}) - 2\partial_\rho \xi^{(\nu} \cdot \gamma^{\mu)\rho*} - c \gamma^{\mu\nu*} \\ \gamma \xi_\mu^* &= \partial_\nu (\xi^\nu \xi_\mu^*) + \xi_\nu^* \partial_\mu \xi^\nu + C_a^* \partial_\mu C^a + c^* \partial_\mu c \\ \gamma c^* &= \partial_\mu (\xi^\mu c^*) \\ \gamma C_a^* &= \partial_\mu (\xi^\mu C_a^*). \end{aligned} \quad (2.6)$$

The cohomological problem to be studied is defined on local functionals. These are local functions integrated over the world-sheet,

$$W^g = \int d^2\sigma w, \quad \text{gh}(w) = g \quad (2.7)$$

where g denotes the ghost number (gh) and the integrands are required to depend polynomially on derivatives of all the fields and antifields. A restriction on the number or order of derivatives that may occur is however *not* imposed. The dependence on the undifferentiated fields X^M and $\gamma_{\mu\nu}$ may of course be nonpolynomial. By definition, a local functional (2.7) is called BRST invariant if the BRST transformation of its integrand is a total derivative,

$$s w = \partial_\mu v^\mu, \quad (2.8)$$

where the v^μ are required to be local functions too. Solutions of the form $s\hat{w} + \partial_\mu \hat{v}^\mu$ are called trivial whenever $\hat{w}$ and $\hat{v}^\mu$ are local functions, and two solutions are called equivalent ($\simeq$) if they differ by a trivial solution,

$$w \simeq w' \quad :\Leftrightarrow \quad w - w' = s\hat{w} + \partial_\mu \hat{v}^\mu. \quad (2.9)$$

3 Contracting homotopies and conformal structure

The first part of the cohomological analysis consists in the construction of contracting homotopies which reduce the cohomological problem considerably and do not depend on the specific choice of the classical action (only the knowledge of the gauge transformations is needed). It applies a technique described in [12] which reveals in this case a ‘conformal structure’ of the models under study and reduces the problem to the cohomology in a space of conformal tensor fields and generalized connections. This conformal structure enables us to reduce the analysis further to a “conformally invariant” cohomological problem in a manner that parallels the analysis of sigma models performed in [13]. We shall not repeat in detail the arguments and constructions used in [12] and [13]; rather we shall focus on the basic ideas and describe afterwards their outcome for the models studied here. These ideas are:

1. The formulation of the cohomological problem in the jet bundle of the fields and antifields. Essentially this means that the fields, antifields and all their derivatives are considered as local coordinates of an infinite jet space J^∞ .
2. The mapping of the BRST cohomology in the space of local functionals to the cohomology of

$$\tilde{s} = s + d \tag{3.1}$$

in the space of local total forms on J^∞ (“ $\tilde{s}$ -cohomology” for short). A local total form is simply a sum of local forms with different form degrees and ghost numbers.

3. The construction of a complete set of new local jet coordinates $\{\mathcal{U}^\ell, \mathcal{V}^\ell, \mathcal{W}^i\}$ satisfying

$$\tilde{s}\mathcal{U}^\ell = \mathcal{V}^\ell, \quad \tilde{s}\mathcal{W}^i = \mathcal{R}^i(\mathcal{W}). \tag{3.2}$$

This allows to contract the $\tilde{s}$ -cohomology on local total forms (locally) to the $\tilde{s}$ -cohomology on local total forms constructed solely of the $\mathcal{W}$ ’s.

4. A second very powerful contracting homotopy which eliminates also almost all of the $\mathcal{W}$ ’s and reduces the cohomological problem to the $\tilde{s}$ -cohomology in the space of local total forms with vanishing “conformal weights” constructed of the few remaining $\mathcal{W}$ ’s.

The mapping to the $\tilde{s}$ -cohomology just compresses the analysis of the so-called descent equations following from (2.8). Namely, written in terms of differential forms, (2.8) reads $s\omega_2 + d\omega_1 = 0$ where $\omega_2 = d\sigma^+ d\sigma^- w$, $\omega_1 = d\sigma^\mu \epsilon_{\nu\mu} v^\nu$ (with $\epsilon_{-+} = 1$). By the standard arguments one concludes then the existence of a 0-form ω_0 such that altogether

$$s\omega_2 + d\omega_1 = 0, \quad s\omega_1 + d\omega_0 = 0, \quad s\omega_0 = 0. \tag{3.3}$$

These are the descent equations. They are equivalent to

$$\tilde{s}\omega = 0, \quad \omega = \omega_2 + \omega_1 + \omega_0 . \quad (3.4)$$

When switching from s to $\tilde{s}$, it is natural to introduce a *total degree* (totdeg) which is the sum of the ghost number and the form degree (i.e. $\tilde{s}$ has total degree 1),

$$\text{totdeg} = \text{gh} + \text{formdeg} . \quad (3.5)$$

Hence, the integrands of the BRST invariant local functionals with ghost number g are just the 2-forms contained in the corresponding $\tilde{s}$ -invariant local total forms with total degree $g + 2$.

The crucial properties of jet coordinates satisfying (3.2) are of course that the $\mathcal{U}$'s and $\mathcal{V}$'s form “ $\tilde{s}$ -doublets”, and that $\tilde{s}$ leaves the space of the $\mathcal{W}$'s invariant. The construction of these jet coordinates given below is in fact technically the most involved part of our analysis. The differences to an analogous construction in [13] are (a) the presence of the abelian gauge and ghost fields and their antifields which amounts to the finding of $\mathcal{U}$'s, $\mathcal{V}$'s and $\mathcal{W}$'s corresponding to these extra fields, and (b) the switch from s to $\tilde{s}$. (b) is actually straightforward as the s -transformations turn into $\tilde{s}$ -transformations by the replacement $\xi^\mu \rightarrow \xi^\mu + d\sigma^\mu$. Nevertheless, switching from s to $\tilde{s}$ simplifies some arguments, such as the proof that explicit world-sheet coordinates can be removed from all the cocycles (as they count among the $\mathcal{U}$'s).

In order to construct the $\mathcal{U}$'s, $\mathcal{V}$'s and $\mathcal{W}$'s we first replace the undifferentiated components of the world-sheet metric and of the ghosts by the following new jet coordinates,

$$\begin{aligned} h_{\pm\pm} &= \gamma_{\pm\pm}(\gamma_{+-} + \sqrt{\gamma})^{-1} \\ e &= \sqrt{\gamma} \\ \tilde{\xi}^\pm &= \xi^\pm + d\sigma^\pm + h_{\mp\mp}(\xi^\mp + d\sigma^\mp) \\ \tilde{c} &= c e + \partial_\mu(\xi^\mu e) + d\sigma^\mu \partial_\mu e \\ \tilde{C}^a &= C^a + (\xi^\mu + d\sigma^\mu) A_\mu^a . \end{aligned} \quad (3.6)$$

The $h_{\pm\pm}$ may be viewed as “Beltrami variables”. The $\tilde{s}$ -transformations of the fields read in the new basis:

$$\begin{aligned} \tilde{s} h_{\pm\pm} &= (\partial_\pm - h_{\pm\pm} \partial_\mp + \partial_\mp h_{\pm\pm}) \tilde{\xi}^\mp \\ \tilde{s} e &= \tilde{c} \\ \tilde{s} A_\pm^a &= \partial_\pm \tilde{C}^a \mp (\tilde{\xi}^\mp - h_{\pm\pm} \tilde{\xi}^\pm) F^a \\ \tilde{s} \tilde{\xi}^\pm &= \tilde{\xi}^\pm \partial_\pm \tilde{\xi}^\pm \\ \tilde{s} \tilde{c} &= 0 \\ \tilde{s} \tilde{C}^a &= \tilde{\xi}^+ \tilde{\xi}^- F^a \\ \tilde{s} X^M &= \tilde{\xi}^\mu \mathcal{D}_\mu X^M \end{aligned} \quad (3.7)$$

where $\mathcal{D}_\mu$ are covariant derivatives given explicitly below, and

$$F^a = \frac{1}{2} (1 - h_{++}h_{--})^{-1} \epsilon^{\mu\nu} F_{\mu\nu}^a, \quad F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a. \quad (3.8)$$

Next one constructs new jet coordinates replacing the derivatives of the fields. Taking also the world-sheet coordinates σ^μ and differentials $d\sigma^\mu$ into account (one has $\tilde{s}\sigma^\mu = d\sigma^\mu$), this yields the new jet coordinates with nonnegative total degrees,

$$\begin{aligned} \{\mathcal{U}_0^\ell\} &= \{\sigma^\mu, \partial_+^m \partial_-^n h_{\pm\pm}, \partial_+^m \partial_-^n e, \partial_+^m \partial_-^n A_-^a, \partial_+^m A_+^a : m, n = 0, 1, \dots\} \\ \{\mathcal{V}_1^\ell\} &= \{\tilde{s}\mathcal{U}_0^\ell\} \\ \{\mathcal{W}_1^i\} &= \{\tilde{C}^a, \partial_\pm^m \tilde{\xi}^\pm : m = 0, 1, \dots\} \\ \{\mathcal{W}_0^i\} &= \{\mathcal{D}_+^m \mathcal{D}_-^n X^M, \mathcal{D}_+^m \mathcal{D}_-^n F^a : m, n = 0, 1, \dots\} \end{aligned} \quad (3.9)$$

where subscripts 0 or 1 indicate the total degree, and $\mathcal{D}_\pm$ are commuting covariant derivatives

$$\mathcal{D}_\pm = (1 - h_{++}h_{--})^{-1} \left[\partial_\pm - h_{\pm\pm} \partial_\mp - \sum_{m \geq 0} (H_\mp^m L_m^\mp - h_{\pm\pm} H_\pm^m L_m^\pm) \right]. \quad (3.10)$$

Here

$$H_\pm^m = \frac{1}{(m+1)!} \partial_\pm^{m+1} h_{\mp\mp}$$

and

$$\begin{aligned} L_r^\pm(\mathcal{D}_\pm^m \mathcal{D}_\mp^n X^M) &= \begin{cases} \frac{m!}{(m-r-1)!} \mathcal{D}_\pm^{m-r} \mathcal{D}_\mp^n X^M & \text{for } 0 \leq r < m \\ 0 & \text{for } r \geq m \end{cases} \\ L_r^\pm(\mathcal{D}_\pm^m \mathcal{D}_\mp^n F^a) &= \begin{cases} \frac{(m+1)!}{(m-r)!} \mathcal{D}_\pm^{m-r} \mathcal{D}_\mp^n F^a & \text{for } 0 \leq r \leq m \\ 0 & \text{for } r > m. \end{cases} \end{aligned} \quad (3.11)$$

The explicit form of the $\mathcal{W}_0^i$ in terms of the fields and their derivatives is obtained from (3.10) and (3.11) in a recursive manner. This is possible because the right hand sides of (3.11) do not contain powers of $\mathcal{D}_\pm$ exceeding m . For instance one gets

$$\begin{aligned} \mathcal{D}_+ X^M &= (1 - h_{++}h_{--})^{-1} (\partial_+ - h_{++} \partial_-) X^M \\ \mathcal{D}_- X^M &= (1 - h_{++}h_{--})^{-1} (\partial_- - h_{--} \partial_+) X^M \\ \mathcal{D}_+ \mathcal{D}_- X^M &= (1 - h_{++}h_{--})^{-1} (\partial_+ - \partial_- h_{++}) \mathcal{D}_- X^M \end{aligned} \quad (3.12)$$

where in the last line ∂_- acts both on h_{++} and $\mathcal{D}_- X^M$. It is also easy to verify by induction that all $\mathcal{W}_0^i$ constructed in this way are local functions because only finitely many summands of the infinite sums in (3.10) are actually nonvanishing thanks to (3.11). We will show below that one can choose the representatives of the cohomology classes such that the only $\mathcal{W}_0^i$ that contribute to them are X^M , F^a and the covariant derivatives of X^M listed in (3.12). Hence, the explicit form of all other $\mathcal{W}_0^i$ will

actually not matter later on. Nevertheless the fact that they exist as local functions of the fields is crucial for our arguments.

The $\mathcal{W}_0^i$ can be viewed as conformal tensor fields, and the $\mathcal{D}_\mu$ as conformal covariant derivatives. Indeed, using (3.11) and the identification

$$L_{-1}^\mu \equiv \mathcal{D}_\mu \quad (3.13)$$

one easily checks that the L 's represent on the $\mathcal{W}_0^i$ two copies of the regular part of the Wit algebra,

$$[L_m^\pm, L_n^\pm] = (m - n)L_{m+n}^\pm, \quad [L_m^+, L_n^-] = 0 \quad (m, n = -1, 0, 1, \dots). \quad (3.14)$$

The occurrence of this infinite dimensional algebra reflects the infinite dimension of the conformal group in two dimension and is related to the Weyl invariance which removes the ‘conformal’ degree of freedom contained in $\gamma_{\mu\nu}$. Indeed, in the above approach this degree of freedom is associated with the variable $e = \sqrt{\gamma}$. Together with the Weyl ghost field, e drops completely out of the cohomological analysis, as these fields and all their derivatives form $\tilde{s}$ -doublets. In fact, we could have worked from the beginning with the ‘Beltrami’ fields h_{++} and h_{--} instead of using the three components of $\gamma_{\mu\nu}$ and imposing Weyl invariance. Now, (3.7) suggests to view h_{++} and h_{--} as gauge fields for ‘chiral diffeomorphisms’, as $\tilde{s}h_{++}$ contains $\partial_+\tilde{\xi}^-$ while $\tilde{s}h_{--}$ contains $\partial_-\tilde{\xi}^+$, but there are no ‘gauge fields’ corresponding to $\partial_+\tilde{\xi}^+$ and $\partial_-\tilde{\xi}^-$. As a consequence, the m th order derivatives of h_{++} and h_{--} pair off with the $(m+1)$ th order derivatives of the ξ^μ except for $\partial_+^{m+1}\tilde{\xi}^+$ and $\partial_-^{m+1}\tilde{\xi}^-$. The undifferentiated ghost fields $\tilde{\xi}^\pm$ and their non-paired derivatives $\partial_\pm^{m+1}\tilde{\xi}^\pm$ correspond to the $L_m^\pm$.

From (3.7) it is obvious that the $\mathcal{W}_1^i$ satisfy (3.2). The $\mathcal{W}_0^i$ satisfy (3.2) too as they transform according to

$$\tilde{s}\mathcal{W}_0^i = \sum_{m \geq -1} \frac{1}{(m+1)!} \left(\partial_+^{m+1}\tilde{\xi}^+ \cdot L_m^+ + \partial_-^{m+1}\tilde{\xi}^- \cdot L_m^- \right) \mathcal{W}_0^i. \quad (3.15)$$

Again, these expressions are local as only finitely many summands are nonvanishing thanks to (3.11).

Next we construct appropriate new local jet coordinates replacing one by one the antifields and all their derivatives such that (3.2) holds. For this construction the knowledge of the classical action is not needed, as $\tilde{\gamma}\mathcal{W}^i = R^i(\mathcal{W})$ (with $\tilde{\gamma} = \gamma + d$) implies $\tilde{s}\mathcal{W}^i = \mathcal{R}^i(\mathcal{W})$ in our case as a consequence of $\tilde{s} = \delta + \tilde{\gamma}$ and $\delta\tilde{\gamma} + \tilde{\gamma}\delta = 0$. Furthermore $\tilde{\gamma}$ does not involve the explicit form of the classical action but only the gauge symmetries, cf. section 2. This is actually the reason that one can use the same $\mathcal{U}$'s, $\mathcal{V}$'s and $\mathcal{W}$'s for all actions with the field content and gauge symmetries that we have imposed.

The construction of new jet coordinates replacing the antifields $\gamma^{\mu\nu*}$, ξ_μ^* , c^* and all their derivatives has been given in [13]. Their explicit form will not be needed later on as none of them will contribute nontrivially to the cohomology. We will denote them by $\partial_+^m \partial_-^n e^*$, $\partial_+^m \partial_-^n \tilde{c}^*$, $h_{m,n}^{\pm\pm*}$, $\xi_{\mu,m,n}^*$ ($m, n = 0, 1, \dots$). The $\partial_+^m \partial_-^n \tilde{c}^*$ and $\partial_+^m \partial_-^n e^*$

count among the $\mathcal{U}$'s and $\mathcal{V}$'s respectively (due to $\tilde{s}\tilde{c}^* = e^*$), whereas the $h_{m,n}^{\pm\pm*}$ and $\xi_{\mu m,n}^*$ are $\mathcal{W}$'s. The new jet coordinates replacing the undifferentiated antifields X_M^* , $A_a^{\mu*}$ and C_a^* are

$$\begin{aligned}\hat{X}_M^* &= (1 - h_{++}h_{--})^{-1}X_M^* \\ \hat{A}_a^{\pm*} &= (1 - h_{++}h_{--})^{-1}(A_a^{\pm*} + h_{\mp\mp}A_a^{\mp*}) \\ \hat{C}_a^* &= (1 - h_{++}h_{--})^{-1}C_a^*.\end{aligned}\tag{3.16}$$

The partial derivatives of X_M^* , $A_a^{\mu*}$ and C_a^* are replaced by covariant derivatives of the redefined antifields (3.16). These covariant derivatives are constructed recursively in the same manner as those of X^M and F^a above, where $L_r^\pm$ acts on the covariant derivatives of $\hat{X}_M^*$, $\hat{A}_a^{\mp*}$ and $\hat{C}_a^*$ as on those of F^a in (3.11), while it acts on the covariant derivatives of $\hat{A}_a^{\pm*}$ as on those of X^M . In particular this yields

$$\mathcal{D}_\pm \hat{A}_a^{\pm*} = (1 - h_{++}h_{--})^{-1}(\partial_\pm - \partial_\mp h_{\pm\pm}) \hat{A}_a^{\pm*}.\tag{3.17}$$

Again, the explicit expressions for other covariant derivatives of $\hat{X}_M^*$, $\hat{C}_a^*$ or $\hat{A}_a^{\mu*}$ will not be needed later on as they will contribute only trivially to the cohomology.

$\hat{X}_M^*$, $\hat{C}_a^*$, $\hat{A}_a^{\mu*}$ and their covariant derivatives count among the $\mathcal{W}$'s and we thus get the following set of new jet coordinates replacing the antifields and all their derivatives:

$$\begin{aligned}\{\mathcal{U}_{-2}^\ell\} &= \{\partial_+^m \partial_-^n \tilde{c}^* : m, n = 0, 1, \dots\} \\ \{\mathcal{V}_{-1}^\ell\} &= \{\partial_+^m \partial_-^n e^* : m, n = 0, 1, \dots\} \\ \{\mathcal{W}_{-1}^i\} &= \{h_{m,n}^{\pm\pm*}, \mathcal{D}_+^m \mathcal{D}_-^n \hat{X}_M^*, \mathcal{D}_+^m \mathcal{D}_-^n \hat{A}_a^{\mu*} : m, n = 0, 1, \dots\} \\ \{\mathcal{W}_{-2}^i\} &= \{\xi_{\pm m,n}^*, \mathcal{D}_+^m \mathcal{D}_-^n \hat{C}_a^* : m, n = 0, 1, \dots\}.\end{aligned}\tag{3.18}$$

As mentioned above, the $\mathcal{W}_{-1}^i$ and $\mathcal{W}_{-2}^i$ satisfy (3.2) as a consequence of their $\tilde{\gamma}$ -transformations. The $\tilde{\gamma}$ -transformations of $\hat{X}_M^*$, $\hat{A}_a^{\mu*}$, $\hat{C}_a^*$ and their covariant derivatives take the same form as the $\tilde{s}$ -transformations of the $\mathcal{W}_0^i$ given in (3.15).

This completes the construction of the $\mathcal{U}$'s, $\mathcal{V}$'s and $\mathcal{W}$'s. As mentioned in the beginning, the construction performed so far allows us now to eliminate all the $\mathcal{U}$'s and $\mathcal{V}$'s from the cohomology (at least locally) and we are left with the $\mathcal{W}$'s. Next we show that actually only very few $\mathcal{W}$'s contribute nontrivially to the cohomology. To that end we extend the definition of L_0^+ and L_0^- to all $\mathcal{W}$'s by

$$L_0^\pm = \left\{ \tilde{s}, \frac{\partial}{\partial(\partial_\pm \tilde{\xi}^\pm)} \right\}.\tag{3.19}$$

This reproduces the action of $L_0^\pm$ in (3.11) and extends it such that all $\mathcal{W}$'s are eigenfunctions of L_0^+ and L_0^- . Indeed one has

$$L_0^\pm \mathcal{W}^i = w^\pm(i) \mathcal{W}^i\tag{3.20}$$

where the eigenvalues $w^\pm(i)$ are integer “conformal weights” listed in table 1.

Fields			Antifields		
totdeg	$\mathcal{W}^i$	$w^+(i), w^-(i)$	totdeg	$\mathcal{W}^i$	$w^+(i), w^-(i)$
1	$\partial_+^m \tilde{\xi}^+$	$m-1, 0$	-2	$\xi_{+m,n}^*$	$m+2, 0$
	$\partial_-^m \tilde{\xi}^-$	$0, m-1$		$\xi_{-m,n}^*$	$0, m+2$
	$\tilde{C}^a$	$0, 0$		$\mathcal{D}_+^m \mathcal{D}_-^n \hat{C}_a^*$	$m+1, n+1$
0	$\mathcal{D}_+^m \mathcal{D}_-^n X^M$	m, n	-1	$\mathcal{D}_+^m \mathcal{D}_-^n \hat{X}_M^*$	$m+1, n+1$
				$h_{m,n}^{++*}$	$0, n+2$
				$h_{m,n}^{--*}$	$m+2, 0$
	$\mathcal{D}_+^m \mathcal{D}_-^n F^a$	$m+1, n+1$		$\mathcal{D}_+^m \mathcal{D}_-^n \hat{A}_a^{+*}$	$m, n+1$
				$\mathcal{D}_+^m \mathcal{D}_-^n \hat{A}_a^{-*}$	$m+1, n$

Table 1: Total degrees and conformal weights of the $\mathcal{W}$'s

As L_0^+ and L_0^- are anticommutators with $\tilde{s}$, they establish contracting homotopies for the $\tilde{s}$ -cohomology in the space of local total forms constructed of the $\mathcal{W}$'s, i.e., this cohomology can be nontrivial only in the intersection of the kernels of L_0^+ and L_0^- . Hence, all nontrivial representatives of the $\tilde{s}$ -cohomology can be brought to the form $\omega(\mathcal{W})$ where $\omega(\mathcal{W})$ has vanishing conformal weights,

$$\tilde{s} \alpha(\mathcal{W}) = 0 \Rightarrow \alpha(\mathcal{W}) = \omega(\mathcal{W}) + \tilde{s} \beta(\mathcal{W}), \quad L_0^\pm \omega = 0. \quad (3.21)$$

It is in fact easy to construct the most general local $\omega(\mathcal{W})$ with vanishing conformal weights. Indeed, table 1 shows that all $\mathcal{W}$'s have a nonnegative conformal weight w^+ or w^- , except for $\tilde{\xi}^+$ and $\tilde{\xi}^-$ which have weights $(-1, 0)$ and $(0, -1)$ respectively. $\tilde{\xi}^+$ and $\tilde{\xi}^-$ are thus the only $\mathcal{W}$'s which can compensate for positive conformal weights of other $\mathcal{W}$'s. As $\tilde{\xi}^+$ and $\tilde{\xi}^-$ are Grassmann odd, each of them can appear at most once in a monomial of $\mathcal{W}$'s. Hence, a $\mathcal{W}^i$ with $w^+(i) > 1$ or $w^-(i) > 1$ cannot appear in a *local* total form with vanishing conformal weights. ω can thus only depend on the undifferentiated $\tilde{\xi}^\mu$ and on those $\mathcal{W}$'s with conformal weights $(0, 0)$, $(1, 0)$, $(0, 1)$ or $(1, 1)$ respectively. Furthermore the $\mathcal{W}$'s with conformal weights $(1, 0)$, $(0, 1)$ or $(1, 1)$ appear necessarily together with $\tilde{\xi}^+$, $\tilde{\xi}^-$ and $\tilde{\xi}^+ \tilde{\xi}^-$ respectively. We conclude that ω can be constructed solely out of the following quantities:

$$\begin{aligned}
\{z^s\} &= \{X^M, \tilde{\Theta}_+, \tilde{\Theta}_-, \tilde{C}^a\} \\
\{y_\pm^A\} &= \{\tilde{X}_\pm^M, \tilde{Y}_\pm, \tilde{A}_{a\pm}^{\mp*}\} \\
\{y_{+-}^\Delta\} &= \{\tilde{X}_{+-}^M, \tilde{F}_{+-}^a, \tilde{X}_{M+-}^*, \tilde{A}_{a+-}^{\mu*}, \tilde{C}_{a+-}^*\}
\end{aligned} \quad (3.22)$$

where

$$\tilde{\Theta}_\pm = \partial_\pm \tilde{\xi}^\pm,$$

$$\begin{aligned}
\tilde{X}_{\pm}^M &= \tilde{\xi}^{\pm} \mathcal{D}_{\pm} X^M, \quad \tilde{Y}_{\pm} = \tilde{\xi}^{\pm} \partial_{\pm}^2 \tilde{\xi}^{\pm}, \quad \tilde{A}_{a\pm}^{\mp*} = \tilde{\xi}^{\pm} \hat{A}_a^{\mp*}, \\
\tilde{X}_{+-}^M &= \tilde{\xi}^+ \tilde{\xi}^- \mathcal{D}_+ \mathcal{D}_- X^M, \quad \tilde{F}_{+-}^a = \tilde{\xi}^+ \tilde{\xi}^- F^a, \\
\tilde{X}_{M+-}^* &= \tilde{\xi}^+ \tilde{\xi}^- \hat{X}_{M+}^*, \quad \tilde{A}_{a+-}^{\pm*} = \tilde{\xi}^+ \tilde{\xi}^- \mathcal{D}_{\pm} \hat{A}_a^{\pm*}, \quad \tilde{C}_{a+-}^* = \tilde{\xi}^+ \tilde{\xi}^- \hat{C}_a^*. \quad (3.23)
\end{aligned}$$

$\{z^s\}$ contains all $\mathcal{W}$'s with conformal weights $(0,0)$, $\{y_+^A\}$ ($\{y_-^A\}$) contains all $\mathcal{W}$'s with conformal weights $(1,0)$ ($(0,1)$) multiplied by $\tilde{\xi}^+$ ($\tilde{\xi}^-$) in order to compensate for their conformal weight, and $\{y_{+-}^{\Delta}\}$ contains all $\mathcal{W}$'s with conformal weights $(1,1)$ multiplied by $\tilde{\xi}^+ \tilde{\xi}^-$. Note that the subscripts $+$ or $-$ of the y 's indicate that they contain $\tilde{\xi}^+$ or $\tilde{\xi}^-$. As the $\tilde{\xi}^{\mu}$ anticommute, the product of two y 's vanishes whenever they have a subscript $+$ or $-$ in common. This allows us to write the most general local $\omega(\mathcal{W})$ with vanishing conformal weights in the form

$$\omega = \mathcal{F}(z) + y_+^A \mathcal{G}_A^+(z) + y_-^A \mathcal{G}_A^-(z) + y_{+-}^{\Delta} \mathcal{H}_{\Delta}(z) + y_+^A y_-^B \mathcal{K}_{AB}(z) \quad (3.24)$$

where $\mathcal{F}(z)$, $\mathcal{G}_A^{\pm}(z)$, $\mathcal{H}_{\Delta}(z)$, $\mathcal{K}_{AB}(z)$ are polynomials in the Grassmann odd quantities $\tilde{\Theta}_+$, $\tilde{\Theta}_-$ and $\tilde{C}^a$ with X -dependent coefficient functions. The remainder of the cohomological analysis is simply a direct analysis of the cocycle condition $\tilde{s}\omega = 0$ with ω as in (3.24), modulo corresponding coboundary terms. To perform it we need to determine first the classical action in order to complete the $\tilde{s}$ -transformations of $\tilde{X}_{M+-}^*$, $\tilde{A}_{a\pm}^{\mp*}$ and $\tilde{A}_{a+-}^{\mu*}$.

4 Action

The classical action has vanishing ghost number and does not depend on antifields. The requirement that the action be gauge invariant is in our case equivalent to its BRST invariance (up to surface terms, of course) as the gauge algebra is closed. The most general gauge invariant local action for the models under study is therefore obtained from the $\tilde{s}$ -cohomology at total degree 2 in the space of antifield independent local total forms. We will now compute this cohomology group and then extract from the result the most general local action with the required properties.

Using the results of the previous section this is actually an easy exercise. We will nevertheless describe it in some detail as the procedure applied here will be also used later in the more compact notation (3.22) to complete the cohomological analysis. From the analysis of the previous section we know that we can restrict ourselves to antifield independent local total forms (3.24) in order to classify the gauge invariant local action functionals. Such total forms thus depend only on the quantities listed in table 2. The $\tilde{s}$ -transformations given in the table are obtained from the formulae of the previous section.

Q	X^M	$\tilde{X}_{\pm}^M$	$\tilde{\Theta}_{\pm}$	$\tilde{C}^a$	$\tilde{X}_{+-}^M$	$\tilde{Y}_{\pm}$	$\tilde{F}_{+-}^a$
$\text{totdeg}(Q)$	0	1	1	1	2	2	2
$\tilde{s}Q$	$\tilde{X}_+^M + \tilde{X}_-^M$	$\mp \tilde{X}_{+-}^M$	$\tilde{Y}_{\pm}$	$\tilde{F}_{+-}^a$	0	0	0

Table 2: Properties of the antifield independent z 's and y 's

The most general antifield independent total form (3.24) with total degree 2 is

$$\begin{aligned}\omega^2 = & \tilde{X}_+^M \tilde{X}_-^N f_{MN}(X) + \tilde{\Theta}_+ \tilde{\Theta}_- f(X) + \tilde{C}^a \tilde{C}^b f_{ab}(X) \\ & + \tilde{X}_\mu^M \tilde{\Theta}_\nu f_M^{\mu\nu}(X) + \tilde{X}_\mu^M \tilde{C}^a f_{Ma}^\mu(X) + \tilde{\Theta}_\mu \tilde{C}^a f_a^\mu(X) \\ & + \tilde{Y}_\mu f^\mu(X) + \tilde{F}_{+-}^a f_a(X) + \tilde{X}_{+-}^M f_M(X)\end{aligned}$$

where the f 's are arbitrary functions of the X 's. Using table 2, one easily computes $\tilde{s}\omega^2$ and derives that the cocycle condition $\tilde{s}\omega^2 = 0$ imposes the following restrictions on the target space functions occurring in ω^2 :

$$\tilde{s}\omega^2 = 0 \Leftrightarrow f = f_{ab} = f_a^\mu = 0, \quad f_M^{\nu\mu} = \partial_M f^\mu, \quad f_{Ma}^\mu = \partial_M h_a \quad (4.1)$$

$$\begin{aligned}\Leftrightarrow \omega^2 = & \tilde{X}_+^M \tilde{X}_-^N f_{MN}(X) + (\tilde{X}_+^M + \tilde{X}_-^M)[\tilde{\Theta}_\mu \partial_M f^\mu(X) + \tilde{C}^a \partial_M h_a(X)] \\ & + \tilde{Y}_\mu f^\mu(X) + \tilde{X}_{+-}^M f_M(X) + \tilde{F}_{+-}^a f_a(X)\end{aligned} \quad (4.2)$$

for some functions $h_a(X)$. Here and throughout the paper we use the notation

$$\partial_M \equiv \frac{\partial}{\partial X^M}.$$

In order to remove those pieces from ω^2 which are $\tilde{s}$ -exact in the space of antifield independent local total forms, we consider next the general antifield independent total form (3.24) with total degree 1,

$$\omega^1 = \tilde{X}_\mu^M d_M^\mu(X) + \tilde{\Theta}_\mu d^\mu(X) + \tilde{C}^a d_a(X).$$

Using table 2 again, one gets

$$\begin{aligned}\tilde{s}\omega^1 = & \tilde{X}_+^M \tilde{X}_-^N [\partial_M d_N^-(X) - \partial_N d_M^+(X)] \\ & + (\tilde{X}_+^M + \tilde{X}_-^M)[\tilde{\Theta}_\mu \partial_M d^\mu(X) + \tilde{C}^a \partial_M d_a(X)] \\ & + \tilde{Y}_\mu d^\mu(X) + \tilde{X}_{+-}^M [d_M^-(X) - d_M^+(X)] + \tilde{F}_{+-}^a d_a(X).\end{aligned} \quad (4.3)$$

From (4.2) and (4.3) we see that by subtracting the $\tilde{s}$ -exact piece $\tilde{s}\omega^1$ from an $\tilde{s}$ -cocycle ω^2 , we can shift some of the target space functions occurring in this cocycle,

$$\omega^2 \rightarrow \omega^2 - \tilde{s}\omega^1 \Leftrightarrow \begin{cases} f^\mu & \rightarrow f^\mu - d^\mu \\ f_M & \rightarrow f_M - (d_M^- - d_M^+) \\ \partial_M h_a & \rightarrow \partial_M (h_a - d_a) \\ f_a & \rightarrow f_a - d_a \\ f_{MN} & \rightarrow f_{MN} - (\partial_M d_N^- - \partial_N d_M^+). \end{cases} \quad (4.4)$$

This shows that we can always remove f^μ , f_M and either h_a or f_a from ω^2 (but, in general, not both h_a and f_a) by subtracting an $\tilde{s}$ -exact local total form from ω^2 . We choose

$$d^\mu(X) = f^\mu(X), \quad d_M^-(X) - d_M^+(X) = f_M(X), \quad \partial_M d_a(X) = \partial_M h_a(X). \quad (4.5)$$

Note that this fixes d^μ , d_M^μ and d_a up to arbitrary functions $e_M = \frac{1}{2}(d_M^- + d_M^+)$ and constant contributions e_a to the d_a . The choice (4.5) shifts also the remaining functions f_{MN} and f_a according to (4.4). We denote these remaining (shifted) functions by G_{MN} , B_{MN} and D_a respectively, where G_{MN} and B_{MN} are the symmetric and antisymmetric part of the shifted f_{MN} respectively. We conclude that, up to trivial ($\tilde{s}$ -exact) contributions, any local and antifield independent $\tilde{s}$ -cocycle with total degree 2 can be chosen to be of the form

$$\omega^2 = \tilde{X}_+^M \tilde{X}_-^N [G_{MN}(X) + B_{MN}(X)] + \tilde{F}_{+-}^a D_a(X) \quad (4.6)$$

where G_{MN} and B_{MN} are symmetric and antisymmetric in their indices respectively, and we still have the freedom to shift B_{MN} and D_a according to

$$\begin{aligned} B_{MN}(X) &\rightarrow B_{MN}(X) - \partial_{[M} e_{N]}(X), \\ D_a(X) &\rightarrow D_a(X) - e_a \end{aligned} \quad (4.7)$$

without changing the cohomological class of ω^2 .

The 2-form $\omega_2 = d\sigma^+ d\sigma^- L_0$ contained in ω^2 is now easily extracted from it,

$$L_0 = (1 - h_{++} h_{--}) [G_{MN}(X) + B_{MN}(X)] \mathcal{D}_+ X^M \cdot \mathcal{D}_- X^N + \frac{1}{2} \epsilon^{\mu\nu} D_a(X) F_{\mu\nu}^a. \quad (4.8)$$

We conclude that the integrand of any action with the required properties is, up to a total derivative, of the form (4.8). In terms of the original fields it reads

$$\begin{aligned} L_0 &= \frac{1}{2} \sqrt{\gamma} \gamma^{\mu\nu} G_{MN}(X) \partial_\mu X^M \cdot \partial_\nu X^N \\ &\quad + \frac{1}{2} \epsilon^{\mu\nu} [B_{MN}(X) \partial_\mu X^M \cdot \partial_\nu X^N + D_a(X) F_{\mu\nu}^a]. \end{aligned} \quad (4.9)$$

It is easy to verify that the redefinitions (4.7) indeed change L_0 only by a total derivative.

Recall that we have derived the above result only by fixing the field content and by requiring locality and gauge invariance under world-sheet diffeomorphisms, local Weyl transformations of the world-sheet metric, and abelian gauge transformations of the A_μ^a . It is of course obvious that any action with a Lagrangian (4.9) has these properties. The nontrivial result we have derived here is that this form of the Lagrangian is unique up to total derivatives.

The cohomological analysis carried out in the remainder of this paper will be performed without specifying the functions $G_{MN}(X)$, $B_{MN}(X)$ or $D_a(X)$, i.e. it applies to almost any model with a Lagrangian of the form (4.9). We only have to exclude special choices of $G_{MN}(X)$, $B_{MN}(X)$ and $D_a(X)$ yielding models with even more gauge symmetries than we have imposed. For such models our approach would not be sufficient because one would have to add extra ghost fields and their antifields corresponding to the additional gauge symmetries. We shall discuss models which are excluded for this reason in appendix A.

Apart from requiring the absence of extra gauge symmetries our analysis of models with a Lagrangian of the form (4.9) will be truly general, i.e. we will not impose further restrictions on G_{MN} , B_{MN} or D_a . In particular we do not assume that $G_{MN}(X)$ is invertible, i.e. we do not assume that it can be viewed as a metric of a target space of all X^M . Among others this allows us to include bosonic (effective) D-string actions of the Born–Infeld type. Indeed, such actions arise from (4.9) for specific choices of G_{MN} , B_{MN} or D_a with degenerate G_{MN} [14]. To give an example, we single out one target space coordinate φ and use the notation

$$X^0 = \varphi, \quad X^M = x^m \quad \text{for} \quad m = M > 0, \quad (4.10)$$

and choose

$$\begin{aligned} G_{M0}(X) &= 0, \quad G_{mn}(X) = (1 - \varphi^2)^{-1/2} e^{-\phi(x)} g_{mn}(x) \\ B_{m0} &= 0, \quad B_{mn}(X) = -\varphi(1 - \varphi^2)^{-1/2} e^{-\phi(x)} b_{mn}(x) + c_{mn}(x) \\ D_a(X) &= -\varphi(1 - \varphi^2)^{-1/2} e^{-\phi(x)} d_a(x) + c_a(x). \end{aligned}$$

Eliminating the auxiliary fields φ and $\gamma_{\mu\nu}$, the Lagrangian (4.9) turns indeed into the Born–Infeld type Lagrangian

$$\begin{aligned} L_{BI} &= e^{-\phi(x)} \sqrt{-\det(\mathcal{G}_{\mu\nu} + \mathcal{F}_{\mu\nu})} + \frac{1}{2} \epsilon^{\mu\nu} [c_a(x) F_{\mu\nu}^a + c_{mn}(x) \partial_\mu x^m \partial_\nu x^n] \\ \mathcal{G}_{\mu\nu} &= g_{mn}(x) \partial_\mu x^m \partial_\nu x^n \\ \mathcal{F}_{\mu\nu} &= d_a(x) F_{\mu\nu}^a + b_{mn}(x) \partial_\mu x^m \partial_\nu x^n. \end{aligned} \quad (4.11)$$

Of course, the ‘dilaton factor’ $\exp(-\phi(x))$ can be absorbed into $g_{mn}(x)$, $b_{mn}(x)$ and $d_a(x)$, analogously to the switch from the ‘string frame’ to the ‘Einstein frame’.

There are further potentially interesting actions of this type associated with different choices of G_{MN} , B_{MN} and D_a . For example, let us consider (4.10) and

$$\begin{aligned} G_{M0}(X) &= 0, \quad G_{mn}(X) = (1 + \varphi^2) g_{mn}(x) \\ B_{m0} &= 0, \quad B_{mn}(X) = 2\varphi b_{mn}(x), \quad D_a(X) = 2\varphi d_a(x). \end{aligned}$$

If we eliminate the auxiliary fields φ and $\gamma_{\mu\nu}$ we get the Lagrangian

$$L = \frac{-\det(\mathcal{G}_{\mu\nu} + \mathcal{F}_{\mu\nu})}{\sqrt{-\det(\mathcal{G}_{\mu\nu})}}. \quad (4.12)$$

5 Result

From (4.9) one obtains the Koszul–Tate part of the BRST transformations of the antifields ϕ_i^* in (2.5),

$$\begin{aligned} \delta X_M^* &= -G_{MN} \partial_\mu (\sqrt{\gamma} \gamma^{\mu\nu} \partial_\nu X^N) - \sqrt{\gamma} \gamma^{\mu\nu} \Gamma_{KLM} \partial_\mu X^K \cdot \partial_\nu X^L \\ &\quad + \frac{1}{2} \epsilon^{\mu\nu} (F_{\mu\nu}^a \partial_M D_a + H_{KLM} \partial_\mu X^K \cdot \partial_\nu X^L) \\ \delta A_a^{\mu*} &= \epsilon^{\mu\nu} \partial_\nu D_a \\ \delta \gamma^{\mu\nu*} &= \sqrt{\gamma} G_{MN} \left(\frac{1}{4} \gamma^{\mu\nu} \gamma^{\rho\sigma} \partial_\rho X^M \cdot \partial_\sigma X^N - \frac{1}{2} \partial^\mu X^M \cdot \partial^\nu X^N \right) \end{aligned} \quad (5.1)$$

with $\partial^\mu \equiv \gamma^{\mu\nu} \partial_\nu$, and

$$\begin{aligned}\Gamma_{KLM} &= \frac{1}{2} (\partial_K G_{LM} + \partial_L G_{KM} - \partial_M G_{KL}), \\ H_{KLM} &= \partial_K B_{LM} + \partial_L B_{MK} + \partial_M B_{KL} .\end{aligned}\tag{5.2}$$

This yields the $\tilde{s}$ -transformations in table 3.

Q	$\text{totdeg}(Q)$	$\tilde{s} Q$
X^M	0	$\tilde{X}_+^M + \tilde{X}_-^M$
$\tilde{\Theta}_\pm$	1	$\tilde{Y}_\pm$
$\tilde{C}^a$	1	$\tilde{F}_{+-}^a$
$\tilde{A}_{a\pm}^{\mp*}$	0	$\pm(\tilde{X}_\pm^M \partial_M D_a - \tilde{A}_{a\pm}^{\mp*})$
$\tilde{X}_\pm^M$	1	$\mp \tilde{X}_{+-}^M$
$\tilde{Y}_\pm$	2	0
$\tilde{C}_{a+-}^{*}$	0	$-\tilde{A}_{a+-}^{+*} - \tilde{A}_{a+-}^{-*}$
$\tilde{A}_{a+-}^{\pm*}$	1	$\pm(\tilde{X}_+^M \tilde{X}_-^N \partial_M \partial_N D_a + \tilde{X}_{+-}^M \partial_M D_a)$
$\tilde{X}_{M+-}^*$	1	$-2G_{MN} \tilde{X}_{+-}^N + (H_{KLM} - 2\Gamma_{KLM}) \tilde{X}_+^K \tilde{X}_-^L$ $+ \tilde{F}_{+-}^a \partial_M D_a$
$\tilde{X}_{+-}^M$	2	0
$\tilde{F}_{+-}^a$	2	0

Table 3: Total degrees and $\tilde{s}$ -transformations of the z 's and y 's

The cohomology can now be computed directly by applying $\tilde{s}$ to the most general local total form (3.24), and elaborating the conditions imposed by $\tilde{s}$ invariance and nontriviality. These are conditions on the functions of the z 's occurring in (3.24), i.e. the aim is to determine these functions. The computation is analogous to the one performed in the previous section to determine the action functionals, but of course more involved. We have relegated it to appendix B and will summarize the results in the following.

In order to present these results we use a “superfield notation”. To this end we expand the functions of the z 's in terms of $\tilde{\Theta}_+$ and $\tilde{\Theta}_-$. As the $\tilde{\Theta}$'s anticommute, this expansion contains only a few terms and is indeed reminiscent of a superfield. Such “superfields” are always denoted by calligraphic capital letters while we use corresponding small latin letters for their “component functions”, such as

$$\mathcal{F}(z) = f(X, \tilde{C}) + \tilde{\Theta}_\mu f^\mu(X, \tilde{C}) + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{f}(X, \tilde{C}).\tag{5.3}$$

Note that the component functions depend on the X^M and $\tilde{C}^a$, i.e. they involve both bosonic and fermionic variables. For notational convenience, we shall often leave out the arguments of the superfields and their component functions.

Using this notation, the result can be described as follows. Modulo $\tilde{s}$ -exact contributions, the most general $\tilde{s}$ -invariant local total form is

$$\begin{aligned}
\omega = & \tilde{F}_{+-}^a \mathcal{H}_a + \tilde{X}_+^M \tilde{X}_-^N \mathcal{K}_{MN} + \tilde{X}_+^M \tilde{Y}_- \mathcal{K}_{M-} + \tilde{Y}_+ \tilde{X}_-^M \mathcal{K}_{+M} \\
& + \tilde{Y}_+ \tilde{Y}_- \mathcal{K} + \tilde{Y}_+ (\mathcal{G}^+ + \tilde{A}_{a-}^{+*} \mathcal{K}_+^a) + \tilde{Y}_- (\mathcal{G}^- + \tilde{A}_{a+}^{-*} \mathcal{K}_-^a) \\
& + \tilde{X}_{M+-}^* \mathcal{H}^M + (\tilde{C}_{a+-}^* + \tilde{A}_{a-}^{+*} - \tilde{A}_{a+}^{-*}) \mathcal{H}^a + \tilde{A}_{a+}^{-*} \tilde{A}_{b-}^{+*} \mathcal{K}^{ab} \\
& + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} + \tilde{X}_-^M \tilde{A}_{a+}^{-*}) \mathcal{A}_M^a + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} - \tilde{X}_-^M \tilde{A}_{a+}^{-*}) \mathcal{B}_M^a \\
& + (\tilde{X}_-^M - \tilde{X}_+^M) G_{MN} \mathcal{H}^N + \frac{1}{2} (\tilde{X}_+^M + \tilde{X}_-^M) \mathcal{P}_M + \mathcal{F}.
\end{aligned} \tag{5.4}$$

Here $\mathcal{H}_a$ and $\mathcal{K}_{MN}$ are arbitrary “superfields”, while the other ones have the following expansions in component functions,

$$\begin{aligned}
\mathcal{K}_{M-} &= \tilde{\Theta}_+ k_M + \tilde{\Theta}_- k_{M-} + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{M-} \\
\mathcal{K}_{+M} &= \tilde{\Theta}_- k_M + \tilde{\Theta}_+ k_{+M} + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{+M} \\
\mathcal{K} &= \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k} \\
\mathcal{K}_\pm^a &= \tilde{\Theta}_\pm k_\pm^{a\pm} \\
\mathcal{G}^\pm &= \tilde{\Theta}_\pm g^{\pm\pm} \\
\mathcal{H}^M &= h^M + \tilde{\Theta}_+ h^{M+} + \tilde{\Theta}_- h^{M-} \\
\mathcal{A}_M^a &= a_M^a + \tilde{\Theta}_+ a_M^{a+} + \tilde{\Theta}_- a_M^{a-} \\
\mathcal{B}_M^a &= b_M^a + \tilde{\Theta}_+ b_M^{a+} + \tilde{\Theta}_- b_M^{a-} \\
\mathcal{P}_M &= p_M + 2G_{MN} (\tilde{\Theta}_- h^{N-} - \tilde{\Theta}_+ h^{N+}) \\
\mathcal{H}^a &= h^a, \quad \mathcal{K}^{ab} = k^{(ab)}, \quad \mathcal{F} = f,
\end{aligned} \tag{5.5}$$

and have to satisfy

$$\partial_{(M} [G_{N)K} \mathcal{H}^K] - \Gamma_{MNK} \mathcal{H}^K + \frac{1}{2} \mathcal{A}_{(M}^a \partial_{N)} D_a = 0 \tag{5.6}$$

$$\partial_{[M} \mathcal{P}_{N]} + H_{MNK} \mathcal{H}^K + \mathcal{B}_{[M}^a \partial_{N]} D_a = 0 \tag{5.7}$$

$$\frac{\partial \mathcal{F}}{\partial \tilde{C}^a} + \mathcal{H}^M \partial_M D_a = 0 \tag{5.8}$$

$$\partial_M \mathcal{F} - \mathcal{H}^a \partial_M D_a = 0 \tag{5.9}$$

$$\partial_M \mathcal{H}^a + \mathcal{K}^{ab} \partial_M D_b = 0 \tag{5.10}$$

$$\partial_M \mathcal{G}^\mp \pm \mathcal{K}_\mp^a \partial_M D_a = 0. \tag{5.11}$$

One can still subtract trivial ($\tilde{s}$ -exact) local total forms $\tilde{s} \hat{\omega}$ from ω without changing the form of ω . These total forms $\hat{\omega}$ are of the form

$$\begin{aligned}
\hat{\omega} = & \hat{f} + \frac{1}{2} (\tilde{X}_+^M + \tilde{X}_-^M) \hat{\mathcal{P}}_M + (\tilde{X}_-^M - \tilde{X}_+^M) [G_{MN} \hat{\mathcal{H}}^N - \hat{\mathcal{Z}}^a \partial_M D_a] \\
& + \tilde{Y}_+ (\hat{g}^+ + \tilde{\Theta}_+ \hat{g}^{++}) + \tilde{Y}_- (\hat{g}^- + \tilde{\Theta}_- \hat{g}^{--})
\end{aligned}$$

$$\begin{aligned}
& +\tilde{X}_{M+-}^*\hat{\mathcal{H}}^M + (\tilde{C}_{a+-}^* + \tilde{A}_{a-}^{+*} - \tilde{A}_{a+}^{-*})\hat{h}^a \\
& +\tilde{X}_+^M\tilde{A}_{a-}^{+*}\hat{\mathcal{K}}_M^a + \tilde{X}_-^M\tilde{A}_{a+}^{-*}\hat{\mathcal{K}}_M^a + \tilde{Y}_+\tilde{A}_{a-}^{+*}\hat{\mathcal{K}}_+^a + \tilde{A}_{a+}^{-*}\tilde{Y}_-\hat{\mathcal{K}}_-^a \\
& +(\tilde{A}_{a+-}^{+*} - \tilde{A}_{a+-}^{-*})\hat{\mathcal{Z}}^a + \tilde{A}_{a+}^{-*}\tilde{A}_{b-}^{+*}\hat{\mathcal{K}}^{ab}
\end{aligned} \tag{5.12}$$

where $\hat{\mathcal{P}}_M$ and $\hat{\mathcal{K}}^{ab}$ are given by

$$\hat{\mathcal{P}}_M = \hat{p}_M + \tilde{\Theta}_\mu \hat{p}_M^\mu + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{\mathcal{P}}_M \tag{5.13}$$

$$\hat{p}_M^\pm = \mp 2G_{MN}\hat{h}^{N\pm} + 2\partial_M \hat{g}^\pm \pm 2(\hat{z}^{a\pm} - \hat{k}_\pm^a)\partial_M D_a \tag{5.14}$$

$$\bar{\mathcal{P}}_M = -(\hat{k}_-^{a+} + \hat{k}_+^{a-})\partial_M D_a \tag{5.15}$$

$$\hat{\mathcal{K}}^{ab} = \hat{k}^{ab} + \tilde{\Theta}_\mu \hat{k}^{ab\mu}. \tag{5.16}$$

In fact, up to $\tilde{s}$ -invariant contributions, $\hat{\omega}$ is the most general local total form such that subtracting $\tilde{s}\hat{\omega}$ from ω only modifies the component functions of the superfields appearing in (5.4) and (5.5), without introducing additional ones. A shift $\omega \rightarrow \omega - \tilde{s}\hat{\omega}$ results in

$$\mathcal{P}_M \rightarrow \mathcal{P}_M - 2\partial_M \hat{f} + 2\hat{h}^a \partial_M D_a \tag{5.17}$$

$$\mathcal{A}_M^a \rightarrow \mathcal{A}_M^a + 2\hat{\mathcal{K}}^{[ab]}\partial_M D_a \tag{5.18}$$

$$\mathcal{B}_M^a \rightarrow \mathcal{B}_M^a - 2\partial_M \hat{h}^a - 2\hat{\mathcal{K}}^{(ab)}\partial_M D_a \tag{5.19}$$

$$\mathcal{H}_a \rightarrow \mathcal{H}_a - \frac{\partial \hat{f}}{\partial \tilde{C}^a} - \hat{\mathcal{H}}^M \partial_M D_a \tag{5.20}$$

$$\begin{aligned}
\mathcal{K}_{MN} & \rightarrow \mathcal{K}_{MN} - (\partial_{[M}\hat{\mathcal{P}}_{N]} + H_{MNK}\hat{\mathcal{H}}^K + \hat{\mathcal{B}}_{[M}^a\partial_{N]}D_a) \\
& - 2(\partial_{(M}[G_{N)K}\hat{\mathcal{H}}^K] - \Gamma_{MNK}\hat{\mathcal{H}}^K + \frac{1}{2}\hat{\mathcal{A}}_{(M}^a\partial_{N)}D_a)
\end{aligned} \tag{5.21}$$

$$k_M \rightarrow k_M + G_{MN}\bar{h}^N - \frac{1}{2}(2\bar{z}^a + \hat{k}_-^{a+} - \hat{k}_+^{a-})\partial_M D_a \tag{5.22}$$

$$k_{M-}^- \rightarrow k_{M-}^- - \hat{k}_-^{a-}\partial_M D_a - \partial_M \hat{g}^{--} \tag{5.23}$$

$$k_{+M}^+ \rightarrow k_{+M}^+ - \hat{k}_+^{a+}\partial_M D_a - \partial_M \hat{g}^{++} \tag{5.24}$$

$$\bar{k}_{+M} \rightarrow \bar{k}_{+M} + \bar{\hat{k}}_+^a\partial_M D_a \tag{5.25}$$

$$\bar{k}_{M-} \rightarrow \bar{k}_{M-} - \bar{\hat{k}}_-^a\partial_M D_a \tag{5.26}$$

where

$$\begin{aligned}
\hat{\mathcal{A}}_M^a & = \hat{\mathcal{K}}_M^a + \hat{\mathcal{K}}_M^a - 2\partial_M \hat{\mathcal{Z}}^a \\
\hat{\mathcal{B}}_M^a & = \hat{\mathcal{K}}_M^a - \hat{\mathcal{K}}_M^a.
\end{aligned} \tag{5.27}$$

All other superfields in ω remain unchanged, i.e. they cannot be removed or modified by subtracting $\tilde{s}$ -exact contributions from ω without changing its form.

(5.4) contains many separately $\tilde{s}$ -invariant solutions. We group them in eight types, three of which appear in two “chiralities”. Among others, this will facilitate

the discussion of the results later on.

$$\omega = \sum_{i=1,2,3,5,6} \omega_i + \sum_{i=4,7,8} (\omega_{i+} + \omega_{i-}), \quad \tilde{s} \omega_i = \tilde{s} \omega_{i\pm} = 0; \quad (5.28)$$

$$\omega_1 = \tilde{F}_{+-}^a \mathcal{H}_a(z) \quad (5.29)$$

$$\omega_2 = \tilde{X}_+^M \tilde{X}_-^N \mathcal{K}_{MN}(z) \quad (5.30)$$

$$\omega_3 = (\tilde{X}_+^M \tilde{Y}_- \tilde{\Theta}_+ + \tilde{Y}_+ \tilde{X}_-^M \tilde{\Theta}_-) k_M(X, \tilde{C}) \quad (5.31)$$

$$\omega_{4+} = \tilde{X}_+^M \tilde{Y}_- [\tilde{\Theta}_- k_{M-}^-(X, \tilde{C}) + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{M-}(X, \tilde{C})] \quad (5.32)$$

$$\omega_{4-} = \tilde{Y}_+ \tilde{X}_-^M [\tilde{\Theta}_+ k_{+M}^+(X, \tilde{C}) + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{+M}(X, \tilde{C})] \quad (5.33)$$

$$\omega_5 = \tilde{Y}_+ \tilde{Y}_- \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}(X, \tilde{C}) \quad (5.34)$$

$$\begin{aligned} \omega_6 = & \tilde{X}_{M+-}^* h^M(X, \tilde{C}) + (\tilde{C}_{a+-}^* + \tilde{A}_{a-}^{+*} - \tilde{A}_{a+}^{-*}) h^a(X, \tilde{C}) \\ & + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} + \tilde{X}_-^M \tilde{A}_{a+}^{-*}) a_M^a(X, \tilde{C}) + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} - \tilde{X}_-^M \tilde{A}_{a+}^{-*}) b_M^a(X, \tilde{C}) \\ & + \tilde{A}_{a+}^{-*} \tilde{A}_{b-}^{+*} k^{(ab)}(X, \tilde{C}) + (\tilde{X}_-^M - \tilde{X}_+^M) G_{MN}(X) h^N(X, \tilde{C}) \\ & + \frac{1}{2} (\tilde{X}_+^M + \tilde{X}_-^M) p_M(X, \tilde{C}) + f(X, \tilde{C}) \end{aligned} \quad (5.35)$$

$$\begin{aligned} \omega_{7\pm} = & -\tilde{\Theta}_\pm \left[\tilde{X}_{M+-}^* h^{M\pm}(X, \tilde{C}) + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} + \tilde{X}_-^M \tilde{A}_{a+}^{-*}) a_M^{a\pm}(X, \tilde{C}) \right. \\ & + \frac{1}{2} (\tilde{X}_+^M \tilde{A}_{a-}^{+*} - \tilde{X}_-^M \tilde{A}_{a+}^{-*}) b_M^{a\pm}(X, \tilde{C}) \\ & \left. \mp 2 \tilde{X}_\pm^M G_{MN}(X) h^{N\pm}(X, \tilde{C}) \right] \end{aligned} \quad (5.36)$$

$$\omega_{8\pm} = \tilde{Y}_\pm \tilde{\Theta}_\pm [g^{\pm\pm}(X, \tilde{C}) + \tilde{A}_{a\mp}^{\pm*} k_{\pm}^{a\pm}(X, \tilde{C})] \quad (5.37)$$

Note that $\omega_1, \dots, \omega_5$ do not depend on antifields. Each of them is $\tilde{s}$ -invariant for any choice of the component functions occurring in them. In contrast, $\omega_6, \omega_{7\pm}$ and $\omega_{8\pm}$ involve in general antifields, and the component functions occurring in them are determined by partial differential equations following from (5.6)–(5.11). Let us now spell out these equations explicitly for the respective solutions.

(5.6)–(5.10) impose the following equations on ω_6 ,

$$\begin{aligned} \partial_{(M} [G_{N)K} h^K] - \Gamma_{MKN} h^K + \frac{1}{2} a_{(M}^a \partial_{N)} D_a &= 0 \\ \partial_{[M} p_{N]} + H_{MKN} h^K + b_{[M}^a \partial_{N]} D_a &= 0 \\ \frac{\partial f}{\partial \tilde{C}^a} + h^M \partial_M D_a &= 0 \\ \partial_M f - h^a \partial_M D_a &= 0 \\ \partial_M h^a + k^{(ab)} \partial_M D_b &= 0 \end{aligned} \quad (5.38)$$

while (5.11) does not constrain the functions in ω_6 . Eqs. (5.38) imply $\tilde{s} \omega_6 = 0$.

(5.6)–(5.8) require the functions in $\omega_{7\pm}$ to solve

$$\partial_M (G_{NK} h^{K\pm}) - (\Gamma_{MKN} \pm \frac{1}{2} H_{MKN}) h^{K\pm} = -(v_M^{a\pm} \partial_N + w_N^{a\pm} \partial_M) D_a$$

$$h^{M\pm}\partial_M D_a = 0$$

$$\text{where } v_M^{a\pm} = \frac{1}{4}(a_M^{a\pm} \mp b_M^{a\pm}), \quad w_M^{a\pm} = \frac{1}{4}(a_M^{a\pm} \pm b_M^{a\pm}). \quad (5.39)$$

In (5.39), we have merged the two equations originating from (5.6) and (5.7) to a single one. (5.9)–(5.11) do not constrain $\omega_{7\pm}$. Eqs. (5.39) imply $\tilde{s}\omega_{7\pm} = 0$.

(5.6)–(5.10) do not constrain $\omega_{8\pm}$. The only conditions imposed by (5.11) on $\omega_{8\pm}$ are

$$\partial_M g^{\pm\pm} = \pm k_{\pm}^{a\pm} \partial_M D_a. \quad (5.40)$$

This implies $\tilde{s}\omega_{8\pm} = 0$.

Let us finally discuss whether ω_6 , $\omega_{7\pm}$ or $\omega_{8\pm}$ provide also antifield independent solutions. Clearly, $\omega_{8\pm}$ does not depend on antifields only if $k_{\pm}^{a\pm} = 0$. (5.40) implies in this case $\partial_M g^{\pm\pm} = 0$, and thus $g^{\pm\pm} = g_0^{\pm\pm}(\tilde{C})$. These solutions are nontrivial because $g^{\pm\pm}$ cannot be redefined by subtracting trivial solutions from ω (it does not appear in (5.17)–(5.26)). Hence, the antifield independent solutions arising from $\omega_{8\pm}$ are

$$\overset{o}{\omega}_{8\pm} = \tilde{Y}_{\pm} \tilde{\Theta}_{\pm} g_0^{\pm\pm}(\tilde{C}). \quad (5.41)$$

ω_6 does not depend on antifields only if $h^M = h^a = a_M^a = b_M^a = k^{(ab)} = 0$. In this case, (5.38) implies $f = \text{constant}$ and $p_M = \partial_M p$ for some $p(X, \tilde{C})$. (5.17) shows that we can assume $\partial_M p = 0$ without loss of generality (by choosing $\hat{f}$ appropriately). Hence, the only nontrivial antifield independent solutions arising from ω_6 are the constants. They can of course be neglected as they do not provide solutions to (2.8).

ω_{7+} and ω_{7-} do not provide antifield independent solutions at all.

6 Discussion of the results

We will now spell out and discuss the physically most important results of our analysis, i.e. the inequivalent solutions of (2.8) with ghost numbers $g \leq 1$. They provide the rigid symmetries, dynamical conservation laws, gauge invariant actions, the first order consistent deformations of these actions and of their gauge symmetries, and the candidate gauge anomalies.

As explained in section 3, the integrands of the nontrivial BRST invariant local functionals with ghost number g are the 2-forms contained in the nontrivial total forms (5.29)–(5.37) with total degree $g + 2$. In order to extract the results at ghost number g , we use in (5.29)–(5.37) expansions in the $\tilde{C}^a$ such as

$$f(X, \tilde{C}) = \sum_{m \geq 0} \frac{1}{m!} \tilde{C}^{a_1} \dots \tilde{C}^{a_m} f_{a_1 \dots a_m}(X). \quad (6.1)$$

Note that we use the same letter f for the ghost independent part $f(X)$ as for the whole function $f(X, \tilde{C})$.

Recall that the antifield dependent total forms (5.35)–(5.37) involve component functions which have to satisfy the partial differential equations (5.38)–(5.40) respectively. Whether or not these equations have nontrivial solutions depends of course

on the particular model, i.e. on the choice of G_{MN} , B_{MN} and D_a in the Lagrangian (4.9). Therefore it also depends on the model whether and which antifield dependent solutions to (2.8) exist. In contrast, the antifield independent solutions do not depend on G_{MN} , B_{MN} and D_a , but notice that it nevertheless depends on the model which of these solutions are nontrivial or inequivalent, as G_{MN} , B_{MN} and D_a enter the cohomologically trivial redefinitions (5.17)–(5.26). In particular, solutions which are equivalent in one model might not be equivalent in another one.

6.1 $g = -2$: Dynamical conservation laws of second order

Nontrivial solutions to (2.8) with ghost number -2 arise solely from the total forms (5.35) with vanishing total degree. These total forms involve only the ghost independent parts of h^a , $k^{(ab)}$ and f . The corresponding solutions to (2.8) are

$$W^{-2} = \int d^2\sigma \left[C_a^* h^a(X) - \frac{1}{2} \epsilon_{\mu\nu} A_a^{\mu*} A_b^{\nu*} k^{(ab)}(X) \right] \quad (6.2)$$

where $h^a(X)$ and $k^{(ab)}(X)$ have to satisfy

$$\partial_M h^a(X) + k^{(ab)}(X) \partial_M D_b(X) = 0. \quad (6.3)$$

(6.3) is obtained from (5.38) and guarantees that W^{-2} is BRST invariant.

Note that $f(X)$ does not occur in (6.2), although it enters the corresponding total form (5.35). The reason is that it contributes only to the 0-form in the descent equations (3.3) corresponding to W^{-2} . (5.38) relates $f(X)$ to $h^a(X)$ through

$$\partial_M f(X) = h^a(X) \partial_M D_a(X). \quad (6.4)$$

As (6.3) guarantees already the BRST invariance of W^{-2} , and as the BRST invariance of W^{-2} implies in turn the descent equations, any solution to (6.3) must imply the existence of a corresponding solution to (6.4). To verify this, we contract (6.3) with $\partial_N D_a$ and antisymmetrize afterwards in M and N . This yields

$$\partial_{[M}(h^a \partial_{N]} D_a) = 0 \Rightarrow \exists f : h^a \partial_M D_a = \partial_M f$$

where we used the ordinary Poincaré lemma for closed 1-forms (on the target space), i.e. $\partial_{[M} V_{N]} = 0 \Rightarrow V_M = \partial_M f$ with $V_M = h^a \partial_M D_a$.

Conversely, any solution to (6.4) implies the existence of a corresponding solution to (6.3). Indeed, differentiating (6.4) with respect to X^N and antisymmetrizing afterwards in M and N yields

$$\partial_{[M} h^a \partial_{N]} D_a = 0 \Rightarrow \exists k^{(ab)} : \partial_M h^a + k^{(ab)} \partial_M D_b = 0. \quad (6.5)$$

Here we have used (A.2)–(A.5), setting there $\eta^M = \lambda^a = a_M^a = 0$ and $b_M^a = \partial_M h^a$.

Hence, (6.3) and (6.4) are actually equivalent conditions. This is in fact not surprising as it illustrates the general feature that the cohomology classes of the local

BRST cohomology at negative ghost numbers $g = -k$ correspond one-to-one to the dynamical local conservation laws of order k [5]. In n dimensions, the latter are represented by weakly (= on-shell) closed $(n - k)$ -forms j_{n-k} ($d j_{n-k} \approx 0$) which are not weakly exact *locally* (in contrast, *topological* conservation laws are locally but not globally d -exact), or, in dual notation, by completely antisymmetric generalized currents $j^{\mu_1 \cdots \mu_k}$ solving

$$\partial_{\mu_1} j^{\mu_1 \cdots \mu_k} \approx 0, \quad j^{\mu_1 \cdots \mu_k} \not\approx \partial_{\mu_0} \hat{j}^{[\mu_0 \cdots \mu_k]}.$$

Now, (6.4) implies that $f(X)$ is a dynamical conservation law of second order ($k = 2$), as it gives

$$d f(X) \approx 0 \Leftrightarrow \partial_\mu j^{\mu\nu} \approx 0, \quad j^{\mu\nu} = \epsilon^{\mu\nu} f(X). \quad (6.6)$$

This is obtained by contracting (6.4) with dX^M due to $dD_a \approx 0$. Note also that $f(X)$ cannot be trivial as it does not contain derivatives (whereas all equations of motions do contain derivatives). We conclude that the nonconstant solutions $f(X)$ to (6.4) exhaust the nontrivial dynamical conservation laws of second order.

It is easy to verify that there are always infinitely many second order conservation laws. Indeed, as we have just mentioned, the functions $D_a(X)$ themselves are conserved. Therefore any function $Z(D(X))$ of the $D_a(X)$ is conserved too. Solutions to (6.3) and (6.4) are thus given by

$$f = Z(D), \quad h^a = \frac{\partial Z}{\partial D_a}, \quad k^{(ab)} = -\frac{\partial^2 Z}{\partial D_a \partial D_b}. \quad (6.7)$$

The solutions W^{-2} themselves have an interpretation too: as explained in [15], they generate rigid symmetries $\Delta_{(2)}$ of the *extended* action $\mathcal{S}$ (= proper solution of the master equation) through $\Delta_{(2)} \cdot = (W^{-2}, \cdot)$. These symmetries act nontrivially only on C^a , A_μ^a and X_M^* according to

$$\begin{aligned} \Delta_{(2)} C^a &= -h^a(X), \quad \Delta_{(2)} A_\mu^a = -\epsilon_{\mu\nu} A_b^{\nu*} k^{(ab)}(X) \\ \Delta_{(2)} X_M^* &= C_a^* \partial_M h^a(X) - \frac{1}{2} \epsilon_{\mu\nu} A_a^{\mu*} A_b^{\nu*} \partial_M k^{(ab)}(X). \end{aligned} \quad (6.8)$$

6.2 $g = -1$: Rigid symmetries and Noether currents

The nontrivial solutions to (2.8) with ghost number -1 arise also solely from (5.35) where this time we have to pick the contribution with total degree 1. The latter contains those pieces of h^a , $k^{(ab)}$ and f which are linear in the $\tilde{C}^a$, i.e. $\tilde{C}^b h_b^a(X)$, $\tilde{C}^c k_c^{(ab)}(X)$ and $\tilde{C}^a f_a(X)$, as well as the ghost independent pieces of h^M , a_M^a , b_M^a and p_M . This yields the following BRST invariant functionals

$$\begin{aligned} W^{-1} &= \int d^2\sigma \left\{ \frac{1}{2} A_a^{\mu*} [b_M^a(X) \partial_\mu X^M - \sqrt{\gamma} \epsilon_{\mu\nu} a_M^a(X) \partial^\nu X^M + 2 A_b^\mu h_b^a(X)] \right. \\ &\quad \left. + X_M^* h^M(X) + C_a^* C^b h_b^a(X) + \frac{1}{2} \epsilon_{\mu\nu} A_a^{\nu*} A_b^{\mu*} C^c k_c^{(ab)}(X) \right\} \quad (6.9) \end{aligned}$$

where $\partial^\mu = \gamma^{\mu\nu} \partial_\nu$ and the target space functions have to solve Eqs. (5.38) which we will discuss now in some more detail. First we note that one of these equations determines $f_a(X)$ in terms of $h^M(X)$ and $D_a(X)$,

$$\frac{\partial f(X, \tilde{C})}{\partial \tilde{C}^a} + h^M(X, \tilde{C}) \partial_M D_a(X) = 0 \Rightarrow f_a(X) = -h^M(X) \partial_M D_a(X). \quad (6.10)$$

The other Eqs. (5.38) give now equations for the remaining target space functions which can be cast in the form

$$\mathcal{L}_h G_{MN}(X) = -a_{(M}^a(X) \partial_{N]} D_a(X) \quad (6.11)$$

$$\mathcal{L}_h B_{MN}(X) = -\partial_{[M} p'_{N]}(X) - b_{[M}^a(X) \partial_{N]} D_a(X) \quad (6.12)$$

$$\mathcal{L}_h \partial_M D_a(X) = -h_a^b(X) \partial_M D_b(X) \quad (6.13)$$

$$\partial_M h_a^b(X) = -k_a^{(bc)}(X) \partial_M D_c(X) \quad (6.14)$$

where $\mathcal{L}_h$ denotes the standard Lie derivative along $h^M(X)$, such as

$$\mathcal{L}_h G_{MN} = h^K \partial_K G_{MN} + \partial_M h^K \cdot G_{KN} + \partial_N h^K \cdot G_{MK},$$

and $p'_M(X)$ is defined by

$$p'_M(X) = p_M(X) + 2B_{MN}(X) h^N(X). \quad (6.15)$$

By the same argument that we used in (6.5), (6.14) is implied by (6.13). Hence, BRST-invariant functionals W^{-1} exist whenever there are solutions to (6.11)–(6.13), and these solutions provide all BRST-invariant local functionals for ghost number -1 up to locally trivial ones. We call equations (6.11)–(6.13) generalized Killing vector equations because (6.11) reduces in the case $D_a = \text{constant}$ to the familiar Killing vector equations, supplemented by the corresponding condition (6.12) discussed in [16, 13].

This implies the results on the rigid symmetries of the models under study announced and discussed in [14]. In particular (6.9) yields the infinitesimal rigid symmetry transformations Δ of the fields which leave the classical action invariant. They are just the coefficient functions of the pieces linear in the antifields of the classical fields:

$$\begin{aligned} \Delta \gamma_{\mu\nu} &= 0, \quad \Delta X^M = h^M(X) \\ \Delta A_\mu^a &= \frac{1}{2} b_M^a(X) \partial_\mu X^M - \frac{1}{2} \sqrt{\gamma} \epsilon_{\mu\nu} a_M^a(X) \partial^\nu X^M + A_\mu^b h_b^a(X). \end{aligned} \quad (6.16)$$

The corresponding Noether currents can be obtained from the antifield independent part of the 1-form contained in (5.35). One finds

$$\begin{aligned} j^\mu &= \sqrt{\gamma} \gamma^{\mu\nu} G_{MN}(X) h^M(X) \partial_\nu X^N \\ &\quad + \epsilon^{\mu\nu} \left[\frac{1}{2} p_M(X) \partial_\nu X^M - A_\nu^a h^M(X) \partial_M D_a(X) \right]. \end{aligned} \quad (6.17)$$

Because of (5.17)–(5.19), the solutions to (6.11)–(6.14) are defined only up to redefinitions

$$\begin{aligned} a_M^a(X) &\rightarrow a_M^a(X) + 2\hat{k}^{[ab]}(X)\partial_M D_b(X) \\ b_M^a(X) &\rightarrow b_M^a(X) - 2\partial_M \hat{h}^a(X) - 2\hat{k}^{(ab)}(X)\partial_M D_b(X) \\ p_M(X) &\rightarrow p_M(X) - 2\partial_M \hat{f}(X) + 2\hat{h}^a(X)\partial_M D_a(X). \end{aligned} \quad (6.18)$$

These redefinitions modify W^{-1} by BRST-exact pieces. Accordingly, they change the rigid symmetry transformations only by special gauge transformations and on-shell trivial symmetries, as we have shown explicitly in [14] to which we also refer for a more detailed discussion of the rigid symmetries. There are models with infinitely many inequivalent rigid symmetries, see [14] and section 7.

6.3 $g = 0$: Gauge invariant actions and continuous consistent deformations

Continuous consistent deformations of a gauge invariant action are modifications of the action and its gauge symmetries, parametrized by deformation parameters, such that the original theory is recovered for vanishing deformation parameters and the deformed action is gauge invariant under the deformed gauge transformations. They are called trivial if they can be removed through local field redefinitions. To first order in the deformation parameters, such deformations are determined by the local BRST cohomology at ghost number 0 [6]. It should be mentioned however that the existence of such deformations might get obstructed at higher orders in the deformation parameters through the BRST cohomology at ghost number 1. Therefore, in general only a subset of nontrivial solutions to (2.8) with ghost number 0 will eventually yield consistent deformations. For this reason we shall speak about possible deformations here.

One may distinguish two types of consistent deformations. First, there are those deformations which do not change the gauge transformations. They add terms to the action which are invariant under the original gauge transformations. In our case (closed gauge algebra) these deformations arise from the antifield independent solutions to (2.8) at ghost number 0, and therefore reproduce the results derived in section 4. The deformations of the second type change both the action and the gauge transformations nontrivially. They arise from solutions to (2.8) with ghost number 0 which depend nontrivially on antifields. The antifield dependent terms in these solutions provide the possible deformations of the gauge transformations and of their algebra to first order in the deformation parameters.

All inequivalent deformations of the first type arise from (5.29) and (5.30). They read respectively

$$W_1^0 = \int d^2\sigma \frac{1}{2} \epsilon^{\mu\nu} h_a(X) F_{\mu\nu}^a \quad (6.19)$$

$$W_2^0 = \int d^2\sigma \frac{1}{2} (\sqrt{\gamma} \gamma^{\mu\nu} + \epsilon^{\mu\nu}) k_{MN}(X) \partial_\mu X^M \cdot \partial_\nu X^N. \quad (6.20)$$

These functionals provide deformations which shift G_{MN} , B_{MN} and D_a by arbitrary functions $k_{(MN)}$, $k_{[MN]}$ and h_a respectively. This reproduces the result of section 4. It should be noted that some shifts of this type are still cohomologically trivial, as follows from (5.20) and (5.21). These shifts are generated by mere local field redefinitions

$$\begin{aligned} X^M &\rightarrow X^M + \hat{h}^M(X) \\ A_\mu^a &\rightarrow A_\mu^a + \frac{1}{2}\hat{b}_M^a(X)\partial_\mu X^M - \frac{1}{2}\sqrt{\gamma}\epsilon_{\mu\nu}\hat{a}_M^a(X)\partial^\nu X^M + A_\mu^b\hat{h}_{0b}^a, \end{aligned}$$

where $\hat{h}_{0b}^a$ is the constant part of $\hat{h}_b^a$. Only this part enters here because the nonconstant part would introduce antifield dependent terms (it contributes in (5.19) and thus in the redefinition of b_{bM}^a which appears in the antifield dependent solution W_6^0 given below).

The possible deformations of the second type arise from the total forms (5.35) and (5.36) with total degree 2. (5.35) yields the following solutions with ghost number 0,

$$\begin{aligned} W_6^0 = \int d^2\sigma \left\{ -A_\mu^a j_a^\mu - \frac{1}{2}\epsilon^{\mu\nu}A_\mu^a A_\nu^b h_a^M(X)\partial_M D_b(X) + X_M^* C^a h_a^M(X) \right. \\ \left. + A_a^{\mu*} C^b \left[\frac{1}{2}b_{bM}^a(X)\partial_\mu X^M - \frac{1}{2}\sqrt{\gamma}\epsilon_{\mu\nu}a_{bM}^a(X)\partial^\nu X^M + A_\mu^c h_{cb}^a(X) \right] \right. \\ \left. + \frac{1}{2}C_a^* C^b C^c h_{bc}^a(X) + \frac{1}{4}\epsilon_{\mu\nu}A_a^{*\nu} A_b^{\mu*} C^c C^d k_{cd}^{(ab)}(X) \right\}, \end{aligned} \quad (6.21)$$

where

$$j_a^\mu = \sqrt{\gamma}G_{MN}(X)h_a^M(X)\partial^\mu X^N + \epsilon^{\mu\nu}\left[\frac{1}{2}p_{Ma}(X)\partial_\nu X^M - h_a^M(X)A_\nu^b\partial_M D_b(X)\right]. \quad (6.22)$$

In (6.21) we have used already that (5.38) gives

$$f_{ab}(X) = h_{[a}^M(X)\partial_M D_{b]}(X).$$

In addition (5.38) requires the target space functions appearing in W_6^0 to satisfy

$$0 = h_{(a}^M(X)\partial_M D_{b)}(X) \quad (6.23)$$

$$\mathcal{L}_a G_{MN} = -a_{a(M}^b \partial_{N)} D_b \quad (6.24)$$

$$\mathcal{L}_a B_{MN} = -\partial_{[M} p'_{N]a} - b_{a[M}^b \partial_{N]} D_b \quad (6.25)$$

$$\mathcal{L}_{[a} \partial_M D_{b]} = h_{ab}^c \partial_M D_c \quad (6.26)$$

$$\partial_M h_{ab}^c = -k_{ab}^{(cd)} \partial_M D_d \quad (6.27)$$

where $\mathcal{L}_a$ denotes the Lie derivative along h_a^M , and

$$p'_{Ma} = p_{Ma} + 2B_{MN}h_a^N.$$

W_6^0 is cohomologically nontrivial unless it can be removed through redefinitions analogous to (6.18),

$$\begin{aligned} a_{aM}^b &\rightarrow a_{aM}^b + 2\hat{k}_a^{[bc]}\partial_M D_c \\ b_{aM}^b &\rightarrow b_{aM}^b - 2\partial_M \hat{h}_a^b - 2\hat{k}_a^{(bc)}\partial_M D_c \\ p_{Ma} &\rightarrow p_{Ma} - 2\partial_M \hat{f}_a + 2\hat{h}_a^b \partial_M D_b . \end{aligned} \quad (6.28)$$

Every nontrivial solution to (6.23)–(6.27) solves also (6.11)–(6.14) nontrivially because (6.23) and (6.26) imply $\mathcal{L}_a \partial_M D_b = h_{ab}^c \partial_M D_c$. The converse is not true, i.e. in general (6.11)–(6.14) has nontrivial solutions which do not solve (6.23)–(6.27) and thus do not give rise to solutions W_6^0 . Hence, as (6.11)–(6.14) determine the rigid symmetries, the deformations (6.21) contain a *subset* of rigid symmetries. This subset can be described as follows. Labelling the elements of a basis for all nontrivial rigid symmetries by an index α , the solutions to (6.11)–(6.14) are sets of target space functions

$$\{h_\alpha^M, a_{M\alpha}^a, b_{M\alpha}^a, p_{M\alpha}, h_{a\alpha}^b, k_{a\alpha}^{(bc)}\}. \quad (6.29)$$

The solutions to (6.23)–(6.27) are then linear combinations of the functions (6.29) with constant coefficients λ_a^α ,

$$h_a^M = \lambda_a^\alpha h_\alpha^M, \dots, k_{ab}^{(cd)} = \lambda_{[a}^\alpha k_{b]\alpha}^{(cd)}, \quad (6.30)$$

subject to the following requirement imposed by (6.23),

$$h_\alpha^M \partial_M D_{(a} \lambda_{b)}^\alpha = 0. \quad (6.31)$$

Accordingly, one has

$$j_a^\mu = \lambda_a^\alpha j_\alpha^\mu \quad (6.32)$$

where j_α^μ is the Noether current (6.17) which corresponds to the α th rigid symmetry.

We conclude that deformations arising from W_6^0 gauge rigid symmetries satisfying (6.31) and deform the abelian gauge transformations corresponding to the A_μ^a . The presence of the term in W_6^0 involving C_a^* indicates that the algebra of the deformed gauge transformations will not be abelian anymore whenever $h_{bc}^a \neq 0$. The term quadratic in the $A_a^{\mu*}$ shows that the algebra of the deformed gauge transformations closes off-shell only if $k_{cd}^{(ab)} = 0$.

Some deformations arising from W_6^0 are obvious, namely standard nonabelian extensions of Lagrangians (4.9) where the abelian field strengths turn into nonabelian ones and the matter fields are coupled minimally to the gauge fields (through standard covariant derivatives). Among them there are deformations of particular models which, after elimination of auxiliary fields, reproduce nonabelian Born-Infeld type Lagrangians [17],

$$L_{BI}^{nonabel.} = \text{Tr} \sqrt{-\det(\mathbf{1}\mathcal{G}_{\mu\nu} + \mathbf{F}_{\mu\nu})}. \quad (6.33)$$

Here $\mathbf{F}_{\mu\nu}$ is a Lie algebra valued matrix containing the nonabelian field strengths, $\mathbf{1}$ is the unit matrix, and $\mathcal{G}_{\mu\nu}$ is an induced world-sheet metric as in (4.11).

Details and more general deformations of this type are given in appendix C. (5.36) yields the following solutions with ghost number 0,

$$W_{7\pm}^0 = \int d^2\sigma \left\{ -2\partial_{\pm} h_{\mp\mp} \cdot h^{M\pm}(X) G_{MN}(X) \mathcal{D}_{\pm} X^N + X_M^* \Theta_{\pm} h^{M\pm}(X) \right. \\ \left. + \frac{1}{2} A_a^{\mu*} \Theta_{\pm} [b_M^{a\pm}(X) \partial_{\mu} X^M - \sqrt{\gamma} \epsilon_{\mu\nu} a_M^{a\pm}(X) \partial^{\nu} X^M] \right\} \quad (6.34)$$

with $h_{\mp\mp}$ and $\mathcal{D}_{\pm} X^M$ as in (3.6) and (3.12) respectively, and

$$\Theta_{\pm} = \partial_{\pm} \xi^{\pm} + h_{\mp\mp} \partial_{\pm} \xi^{\mp}. \quad (6.35)$$

The target space functions $h^{M\pm}(X)$, $a_M^{a\pm}(X)$ and $b_M^{a\pm}(X)$ in $W_7^{0\pm}$ have to satisfy the partial differential equations (5.39). $W_7^{0\pm}$ is nontrivial unless it can be removed through the cohomologically trivial redefinitions

$$a_M^{a\pm}(X) \rightarrow a_M^{a\pm}(X) + 2\hat{k}^{[ab]\pm}(X) \partial_M D_b(X) \\ b_M^{a\pm}(X) \rightarrow b_M^{a\pm}(X) - 2\hat{k}^{(ab)\pm}(X) \partial_M D_b(X). \quad (6.36)$$

Every solution to (5.39) is also a solution to (6.11)–(6.14) with $h_a^b = k_a^{(bc)} = 0$. Furthermore, such a solution to (6.11)–(6.14) is nontrivial whenever the solution to (5.39) is nontrivial too¹. Hence, the nontrivial solutions to (5.39) are also a subset of rigid symmetries. The antifield independent term in W_7^{0+} (W_7^{0-}) contains the $-$ ($+$) component of the Noether currents (6.17) corresponding to these special rigid symmetries due to $2h^{M\pm} G_{MN} \mathcal{D}_{\pm} X^N = j^{\mp}$.

As $W_7^{0\pm}$ involves the diffeomorphism ghosts, corresponding deformations modify the world-sheet diffeomorphisms. The algebra of the diffeomorphisms will however not get deformed at first order in the deformation parameters because $W_7^{0\pm}$ contains neither antifields of ghosts nor terms quadratic in antifields. Analogous deformations of the world-sheet diffeomorphisms in standard sigma models have been found and discussed in [18].

Remark:

The existence of a cocycle (6.21) implies that there are infinitely many rigid symmetries. In other words, an action with Lagrangian (4.9) has infinitely many rigid symmetries whenever the set of equations (6.23)–(6.27) has a nontrivial solution. Namely, one easily verifies that the antibracket (W^{-2}, W_6^0) of cocycles (6.2) and (6.21) has precisely the form (6.9) with $h^M(X) = h^a(X) h_a^M(X)$ etc.. Now, the antibracket of two BRST cocycles is again a BRST cocycle. Furthermore, a cocycle (6.9) corresponds to a nontrivial rigid symmetry whenever $h^M(X) \neq 0$. As there are always infinitely many cocycles W^{-2} (see section 6.1), we conclude that any cocycle W_6^0 gives indeed rise to infinitely many rigid symmetries via the antibrackets (W^{-2}, W_6^0) .

¹Indeed, assume the solution to (6.11)–(6.14) arising from a nontrivial solution to (5.39) were trivial. This means $h^{M\pm} = 0$, $a_M^{a\pm} = -2\hat{k}^{[ab]}\partial_M D_b$, $b_M^{a\pm} = 2\partial_M \hat{h}^a + 2\hat{k}^{(ab)}\partial_M D_b$ and $p_M^{\pm} = 2\partial_M \hat{f} - 2\hat{h}^a \partial_M D_a$. Since solutions to (5.39) fulfill $p_M^{\pm} = \mp 2G_{MN} h^{N\pm}$, one gets $\partial_M \hat{f} = \hat{h}^a \partial_M D_a$ which implies $\partial_M \hat{h}^a = \Lambda^{(ab)} \partial_M D_b$ for some $\Lambda^{(ab)}$ by the same arguments used to derive (6.5). Altogether this contradicts that the solution to (5.39) is nontrivial, because we get $h^{M\pm} = 0$, $a_M^{a\pm} = -2\hat{k}^{[ab]}\partial_M D_b$ and $b_M^{a\pm} = 2\hat{k}'^{(ab)}\partial_M D_b$ with $\hat{k}'^{(ab)} = \hat{k}^{(ab)} + \Lambda^{(ab)}$.

6.4 $g = 1$: Candidate gauge anomalies

The candidate gauge anomalies arise from (5.29), (5.30) and (5.34)–(5.36). Those obtained from (5.29) and (5.30) can be subdivided into two subsets involving the abelian ghosts C^a and the diffeomorphism ghosts respectively,

$$W_{1\pm}^1 = \int d^2\sigma \frac{1}{2} \Theta_{\pm} \epsilon^{\mu\nu} F_{\mu\nu}^a h_a^{\pm}(X) \quad (6.37)$$

$$W_{2\pm}^1 = \int d^2\sigma \frac{1}{2} \Theta_{\pm} (\sqrt{\gamma} \gamma^{\mu\nu} + \epsilon^{\mu\nu}) k_{MN}^{\pm}(X) \partial_{\mu} X^M \cdot \partial_{\nu} X^N \quad (6.38)$$

$$W_{1C}^1 = \int d^2\sigma \frac{1}{2} C^a \epsilon^{\mu\nu} F_{\mu\nu}^b h_{ab}(X) \quad (6.39)$$

$$W_{2C}^1 = \int d^2\sigma \frac{1}{2} C^a (\sqrt{\gamma} \gamma^{\mu\nu} + \epsilon^{\mu\nu}) k_{MNa}(X) \partial_{\mu} X^M \cdot \partial_{\nu} X^N \quad (6.40)$$

with $\Theta_{\pm}$ as in (6.35). These functionals are BRST invariant for any choice of the target space functions occurring in them. They are nontrivial except for those which can be removed through redefinitions obtained from (5.20) and (5.21).

Note that the candidate anomalies (6.37) and (6.38) come in two chiralities. The solutions (6.38) are familiar from ordinary sigma models where particular (“left-right symmetric”) linear combinations of such solutions provide the conformal anomalies involving the β -functions corresponding to G_{MN} and B_{MN} , see [13] for details. Therefore one would expect (6.37) and (6.38) to have a similar interpretation in the models under study.

The candidate anomalies (6.39) and (6.40) are of the form “abelian ghost $\times$ Weyl- and $U(1)$ -invariant scalar density”. Such solutions are familiar from Yang–Mills theory and gravity [19]. In particular, (6.39) reproduces the representatives of standard abelian chiral anomalies for the special choice $h_{ab} = h_{(ab)} = \text{constant}$.

(5.35) and (5.36) yield the following candidate gauge anomalies,

$$\begin{aligned} W_6^1 = \int d^2\sigma \Big\{ & -C^a A_{\mu}^b [\sqrt{\gamma} G_{MN}(X) h_{ab}^M(X) \partial^{\mu} X^N + \frac{1}{2} \epsilon^{\mu\nu} p_{Mab}(X) \partial_{\nu} X^M] \\ & - \frac{1}{2} C^{[a} A_{\mu}^b A_{\nu}^c] \epsilon^{\mu\nu} h_{ab}^M(X) \partial_M D_c(X) + \frac{1}{2} X_M^* C^a C^b h_{ab}^M(X) \\ & + \frac{1}{4} A_a^{\mu*} C^b C^c [b_{Mbc}^a(X) \partial_{\mu} X^M - \sqrt{\gamma} \epsilon_{\mu\nu} a_{Mbc}^a(X) \partial^{\nu} X^M + 2 A_{\mu}^d h_{bcd}^a(X)] \\ & + \frac{1}{6} C^a C^b C^c [C_d^* h_{abc}^d(X) + \frac{1}{2} \epsilon_{\mu\nu} A_e^{*\nu} A_d^{\mu*} k_{abc}^{(ed)}(X)] \Big\} \quad (6.41) \end{aligned}$$

$$\begin{aligned} W_{7\pm}^1 = \int d^2\sigma \Big\{ & 2[\Theta_{\pm}(A_{\mp}^a - h_{\mp\mp} A_{\pm}^a) - C^a \partial_{\pm} h_{\mp\mp}] G_{MN}(X) h_a^{M\pm}(X) \mathcal{D}_{\pm} X^N \\ & - \frac{1}{2} C^a \Theta_{\pm} A_b^{\mu*} [b_{Ma}^{b\pm}(X) \partial_{\mu} X^M - \sqrt{\gamma} \epsilon_{\mu\nu} a_{Ma}^{b\pm}(X) \partial^{\nu} X^M] \\ & - C^a \Theta_{\pm} X_M^* h_a^{M\pm}(X) \Big\}, \quad (6.42) \end{aligned}$$

where the target space functions have to satisfy sets of partial differential equations analogous to (6.23)–(6.27) and (5.39) respectively. These solutions are nontrivial

except for those we can be removed through redefinitions analogous to (6.18) and (6.36). Both (6.41) and (6.42) thus involve subsets of rigid symmetries. The antifield independent parts of (6.41) and (6.42) involve the corresponding Noether currents.

Note that solutions (6.41) occur only if there are at least two abelian gauge fields. In this respect they are somewhat similar to by now well-known candidate anomalies in Yang–Mills theory and gravity [20]. (6.42) gives candidate gauge anomalies which involve necessarily both the abelian ghosts C^a and the diffeomorphism ghosts. These solutions might therefore have an interpretation as mixed anomalies for world-sheet diffeomorphisms and $U(1)$ -transformations.

Finally, (5.37) gives the following candidate anomalies

$$W_{8\pm}^1 = \pm \int d^2\sigma \left\{ g^{\pm\pm}(X) [\Theta_{\pm} \partial_{\pm}^2 h_{\mp\mp} - \partial_{\pm} h_{\mp\mp} \cdot (\partial_{\pm} \Theta_{\pm} + \partial_{\pm} h_{\mp\mp} \cdot \partial_{\pm} \xi^{\mp})] \right. \\ \left. - (A_a^{\pm*} + h_{\mp\mp} A_a^{\mp*}) k_{\pm}^{a\pm}(X) \Theta_{\pm} (\partial_{\pm}^2 \xi^{\pm} + h_{\mp\mp} \partial_{\pm}^2 \xi^{\mp} - 2 \partial_{\pm} h_{\mp\mp} \cdot \partial_{\pm} \xi^{\mp}) \right\} \quad (6.43)$$

where

$$\partial_M g^{\pm\pm} \mp k_{\pm}^{a\pm} \partial_M D_a = 0. \quad (6.44)$$

Up to the constant parts of g^{++} and g^{--} , the solutions to (6.44) determine the conservation laws of second order, cf. section 6.1. Hence, these conservation laws enter the candidate anomalies (6.43). The constant parts of $g^{\pm\pm}$ provide antifield independent solutions arising from (5.41). Fixing these constants to $\mp 1/2$ and performing a partial integration, the corresponding solutions read

$$\overset{o}{W}_{8\pm}^1 = \int d^2\sigma (\xi^{\pm} + h_{\mp\mp} \xi^{\pm}) \partial_{\pm}^3 h_{\mp\mp} \quad . \quad (6.45)$$

These solutions are familiar from ordinary sigma models where their left-right symmetric linear combination is cohomologically equivalent to the conformal anomaly $\int d^2\sigma c \sqrt{\gamma} R$ [13].

7 Example

In order to illustrate the general results we shall now specify them for a simple class of models which we treated already in [14]. These models contain only one abelian gauge field A_{μ} and besides are characterized by

$$G_{0M} = 0, \quad G_{mn} = g(\varphi) \eta_{mn} \\ B_{0m} = 0, \quad B_{mn} = B_{mn}(\varphi), \quad D = D(\varphi), \quad (7.1)$$

where we used the same notation as in (4.10). We will carry out the analysis for any choice of $g(\varphi)$, $B_{mn}(\varphi)$ and $D(\varphi)$ except that we assume $D \neq \text{constant}$ (as the last term in (4.9) becomes a total derivative for $D = \text{constant}$).

Since these models contain only one abelian gauge field, nontrivial solutions to (2.8) exist only at ghost numbers ranging from -2 to 5 (in general, section 5 provides

solutions at ghost numbers $-2, \dots, 4 + N$ where N is the number of abelian gauge fields). As the specification of the antifield independent solutions is straightforward, we shall only discuss the antifield dependent ones. This amounts to solving the partial differential equations (5.38)–(5.40) in the various ghost number sectors.

$g = -2$: Second order conservation laws. The solutions to (6.3) are in this case $h^a \equiv h(\varphi)$ and $k^{(ab)} \equiv -h'(\varphi)/D'(\varphi)$, where $h(\varphi)$ is an arbitrary function of φ and the prime denotes differentiation with respect to φ ,

$$' = \frac{\partial}{\partial \varphi}.$$

The nontrivial solutions of (2.8) at ghost number -2 are thus

$$W^{-2} = \int d^2\sigma \left[C^* h(\varphi) + \frac{1}{2} \epsilon_{\mu\nu} A^{\mu*} A^{\nu*} \frac{h'(\varphi)}{D'(\varphi)} \right]. \quad (7.2)$$

The corresponding second order conservation laws are functions $f(\varphi)$ related to $h(\varphi)$ by (6.4), i.e. one has $h = f'/D'$.

$g = -1$: Rigid symmetries and Noether currents. (6.11)–(6.13) have been solved already in [14]. (6.14) just determines $k \equiv k_a^{(bc)}$ in terms of $h \equiv h_a^b$ and D . Altogether, the general solution of (6.11)–(6.14) reads, up to redefinitions (6.18),

$$\begin{aligned} h^0 &= h^0(\varphi) \\ h^m &= -h^0(\varphi) x^m g'(\varphi) / [2g(\varphi)] + r^m(\varphi) + r^{[mn]}(\varphi) \eta_{mk} x^k \\ a_m &= -2\eta_{mn} h^{n'} g / D', \quad a_0 = 0 \\ b_m &= [2B'_{mn} h^n + (B'_{mn} h^0)' x^n] / D', \quad b_0 = 0 \\ p_m &= B'_{mn} h^0 x^n, \quad p_0 = 0 \\ h &= h^{0'} + h^0 D'' / D' \\ k &= -h' / D'. \end{aligned} \quad (7.3)$$

Here h^0 , r^m and $r^{[mn]}$ are arbitrary functions of φ . The corresponding solutions to (2.8), rigid symmetries and Noether currents are obtained by inserting (7.3) in (6.9), (6.16) and (6.17) respectively. As discussed in [14], these infinitely many rigid symmetries constitute a loop version of the Weyl algebra (= Poincaré algebra + dilatations). Thereby h^0 , r^m and $r^{[mn]}$ give rise to generalized dilatations, translations and Lorentz transformations respectively.

$g = 0$: Deformations of the gauge symmetries. The antifield dependent solutions of (2.8) with ghost number 0 are determined by Eqs. (6.23)–(6.27) and (5.39) respectively. As explained in section 6.3, the nontrivial solutions to (6.23)–(6.27) form a subset of the nontrivial solutions to (6.11)–(6.14) subject to (6.23). In this

case (6.23) yields $h^M \partial_M D = 0 \Leftrightarrow h^0 = 0$. The nontrivial solutions to (6.23)–(6.27) are thus obtained from (7.3) simply by setting $h^0 = 0$ there,

$$\begin{aligned}
h^0 &= 0 \\
h^m &= r^m(\varphi) + r^{[mn]}(\varphi) \eta_{nk} x^k \\
a_m &= -2\eta_{mn} h^{n'} g / D', \quad a_0 = 0 \\
b_m &= 2B'_{mn} h^n / D', \quad b_0 = 0 \\
p_M &= 0, \quad h = 0, \quad k = 0.
\end{aligned} \tag{7.4}$$

Inserting this in (6.21), we get

$$\begin{aligned}
W_6^0 &= \int d^2\sigma \left\{ \frac{1}{2} A^{\mu*} C [b_m \partial_\mu x^m - \sqrt{\gamma} \epsilon_{\mu\nu} a_m \partial^\nu x^m] \right. \\
&\quad \left. + x_m^* C h^m(X) - \sqrt{\gamma} A_\mu g(\varphi) \eta_{mn} h^m \partial^\mu x^n \right\}
\end{aligned} \tag{7.5}$$

with h^m , a_m and b_m as in (7.4). Note that W_6^0 contains the generalized translations and Lorentz transformations, but no dilatations. Hence, corresponding deformations will not gauge the generalized dilatations.

(5.39) requires in this case in particular $h^{M\pm} \partial_M D = 0$ and $\eta_{nk} \partial_m h^{k\pm} = 0$. It is then straightforward to verify that the general solution to (5.39) is, up to trivial redefinitions (6.36),

$$\begin{aligned}
h^{0\pm} &= 0, \quad h^{m\pm} = r^{m\pm}(\varphi) \\
a_m^\pm &= -2\eta_{mn} r^{n\pm'} g / D', \quad a_0^\pm = 0 \\
b_m^\pm &= 2[B'_{mn} r^{n\pm} \mp \eta_{mn} (g r^{n\pm})'] / D', \quad b_0^\pm = 0,
\end{aligned} \tag{7.6}$$

where $r^{m\pm}(\varphi)$ are arbitrary functions. Hence, only the generalized translations, but neither the dilatations nor the Lorentz transformations enter the corresponding solutions (6.34) which read

$$\begin{aligned}
W_7^{0\pm} &= \int d^2\sigma \left\{ \frac{1}{2} A^{\mu*} \Theta_\pm [b_m^\pm(\varphi) \partial_\mu x^m - \sqrt{\gamma} \epsilon_{\mu\nu} a_m^\pm(\varphi) \partial^\nu x^m] \right. \\
&\quad \left. + x_m^* \Theta_\pm r^{m\pm}(\varphi) - 2\partial_\pm h_{\mp\mp} \cdot \eta_{mn} r^{m\pm}(\varphi) g(\varphi) \mathcal{D}_\pm x^n \right\}.
\end{aligned} \tag{7.7}$$

$g = 1$: Antifield dependent candidate gauge anomalies. Solutions with ghost number 1 depending nontrivially on antifields arise only from (6.42) and (6.43), as (6.41) can give solutions only if there are at least two abelian gauge fields. (6.42) gives

$$\begin{aligned}
W_7^{1\pm} &= \int d^2\sigma \left\{ 2[\Theta_\pm (A_\mp - h_{\mp\mp} A_\pm) - C \partial_\pm h_{\mp\mp}] \eta_{mn} r^{n\pm}(\varphi) g(\varphi) \mathcal{D}_\pm x^m \right. \\
&\quad \left. + x_m^* C \Theta_\pm r^{m\pm}(\varphi) + \frac{1}{2} C \Theta_\pm A^{\mu*} [b_m^\pm(\varphi) \partial_\mu x^m - \sqrt{\gamma} \epsilon_{\mu\nu} a_m^\pm(\varphi) \partial^\nu x^m] \right\}
\end{aligned} \tag{7.8}$$

with $a_m^\pm$ and $b_m^\pm$ as in (7.6).

The solutions to (6.44) are

$$g^{\pm\pm} = g^{\pm\pm}(\varphi), \quad k_{\pm}^{\pm} = \pm g^{\pm\pm'}(\varphi)/D'(\varphi). \quad (7.9)$$

Inserting this into (6.43), one obtains the remaining antifield dependent candidate anomalies.

$g > 1$. Further antifield dependent solutions exist in this case only at ghost number 2. They arise from (5.37) through the $\tilde{C}$ -dependent contributions to $g^{\pm\pm}$ and $k_{\pm}^{\pm}$.

8 Conclusions

Let us finally summarize our main results for models with the field content, gauge symmetries and locality requirements described in section 1 and 2. First, we have *proved* the results announced and discussed already in [14], that is:

a) Any Lagrangian describing such a model can be cast in the form (4.9). The class of models can thus be parametrized through target space functions $G_{MN}(X)$, $B_{MN}(X)$ and $D_a(X)$ where G_{MN} and B_{MN} are symmetric and antisymmetric respectively. Particular choices of these functions yield Born–Infeld actions upon elimination of auxiliary fields. We note that none of our results singles out Born–Infeld actions in the larger class of models studied here. Of course, this does not exclude the existence of other criteria which could distinguish them, such as T duality or the existence of supersymmetric extensions with local kappa symmetry.

b) The nontrivial rigid symmetries and Noether currents of such models are precisely given by (6.16) and (6.17) in terms of the solutions to the generalized Killing vector equations (6.11)–(6.14). These equations can have infinitely many inequivalent solutions, i.e. there are models with infinitely many nontrivial rigid symmetries and Noether currents.

In addition we have derived the following results:

c) Apart from the Noether currents, there are infinitely many dynamical conservation laws of second order and corresponding second order symmetries of the master equation, see section 6.1. The occurrence of an infinite number of such conservation laws originates from the fact that the world-sheet is two dimensional and is familiar from two dimensional Maxwell theory, see [5], section 12. We have also seen that this implies an infinite number of rigid symmetries of the classical action whenever there is at least one nontrivial solution to equations (6.23)–(6.27), see remark at the end of section 6.3.

d) The possible nontrivial consistent deformations of the gauge transformations arise from solutions to Eqs. (6.23)–(6.27) and (5.39). These equations determine subsets of the rigid symmetries respectively. The consistent deformations thus gauge some of the rigid symmetries. The accordingly deformed actions involve the corresponding Noether currents. Deformations arising from solutions to (6.23)–(6.27) deform the abelian gauge transformations. Particular examples for such deformations give rise to standard nonabelian Born–Infeld actions. Other deformations of

this kind might even lead to an open gauge algebra. Deformations involving solutions to (5.39) deform the world-sheet diffeomorphisms nontrivially. It could be interesting to examine whether such deformations can provide manifestly $SL(2, Z)$ invariant formulations of the D-string as in [2, 3].

e) There are three types of antifield independent candidate anomalies. Those of the first type, given by (6.39) and (6.40), involve the abelian ghost fields, those of the second type, given by (6.37) and (6.38), the diffeomorphism ghosts. The former take the form “abelian ghost $\times$ density of a gauge invariant action”, the latter take an analogous form where the abelian ghosts are replaced by (6.35). The third type of antifield independent candidate anomalies contains the two solutions (6.45) which are familiar from ordinary sigma models.

f) In sharp contrast to the result on potential anomalies of ordinary sigma models [13], there may also be candidate anomalies from which the antifield dependence *cannot* be removed. One may distinguish three types of them. Those of the first type, given by (6.41), involve a subset of the rigid symmetries determined by equations analogous to (6.23)–(6.27). They contain the abelian ghosts, but not the diffeomorphism ghosts, and can occur only in presence of at least two abelian gauge fields. Those of the second type, given by (6.42), involve the subset of the rigid symmetries determined by (5.39) and contain both the abelian and the diffeomorphism ghosts. Those of the third type, given by (6.43), involve the second order conservation laws and contain only the diffeomorphism ghosts. To our knowledge, none of these antifield dependent candidate anomalies was previously known for models studied here.

Acknowledgements

JS thanks Fundació Agustí Pedro i Pons for financial support. FB was supported by the Spanish ministry of education and science (MEC). Furthermore this work has been partially supported by AEN95-0590 (CICYT), GRQ93-1047 (CIRIT) and by the Commission of European Communities CHRX93-0362 (04).

A Spurious gauge symmetries

In this appendix we discuss some conditions which have to be imposed on the Lagrangians (4.9) in order to prevent the occurrence of additional gauge symmetries. We shall also argue that these conditions are natural as they just eliminate spurious gauge symmetries.

Assume one would choose $D_a(X)$ such that there are target space functions $k^a(X)$ satisfying $k^a \partial_M D_a = 0$. Then the action would be invariant (up to a surface term) under

$$\delta_\varepsilon \gamma_{\mu\nu} = \delta_\varepsilon X^M = 0, \quad \delta_\varepsilon A_\mu^a = \varepsilon_\mu k^a(X)$$

where the ε_μ are arbitrary fields. Hence, in such a case the action would have extra gauge symmetries and one would have to introduce extra ghost fields C_μ and their

antifields $C^{\mu*}$. Therefore we require in particular that the functions $D_a(X)$ are such that

$$k^a \partial_M D_a = 0 \quad \Rightarrow \quad k^a = 0. \quad (\text{A.1})$$

This condition appears natural. Indeed, assume for instance that each function D_a depends only on one target space coordinate and that all these coordinates are different (in fact, this can be often achieved by an appropriate target space reparametrization). Then (A.1) is always satisfied, provided none of the D_a is constant. As a constant D_a yields only a contribution to the Lagrangian which is a total derivative, the corresponding gauge field A_μ^a actually does not contribute to the classical action, and we can indeed assume (A.1) with no loss of generality in such a case.

More generally we require

$$G_{MN} \eta^N = \lambda^a \partial_M D_a, \quad \eta^M \partial_M D_a = 0 \quad (\text{A.2})$$

$$\partial_{(M} [\lambda^a \partial_{N)} D_a] - \Gamma_{MNK} \eta^K + \frac{1}{2} a_{(M}^a \partial_{N)} D_a = 0 \quad (\text{A.3})$$

$$H_{MNK} \eta^K + b_{[M}^a \partial_{N]} D_a = 0 \quad (\text{A.4})$$

$$\Rightarrow \quad \eta^M = \lambda^a = 0, \quad \exists \Lambda^{ab} : \quad a_M^a = \Lambda^{[ab]} \partial_M D_b, \quad b_M^a = \Lambda^{(ab)} \partial_M D_b. \quad (\text{A.5})$$

Namely, assume there were functions $\eta^M(X)$, $\lambda^a(X)$, $a_M^a(X)$ and $b_M^a(X)$ satisfying Eqs. (A.2)–(A.4). Then the Lagrangian would be invariant up to a total derivative under

$$\begin{aligned} \delta_\varepsilon \gamma_{\mu\nu} &= 0, \quad \delta_\varepsilon X^M = \varepsilon \eta^M(X) \\ \delta_\varepsilon A_\mu^a &= \sqrt{\gamma} \epsilon_{\mu\nu} \lambda^a(X) \partial^\nu \varepsilon + \frac{1}{2} \varepsilon [b_M^a(X) \partial_\mu X^M - \sqrt{\gamma} \epsilon_{\mu\nu} a_M^a(X) \partial^\nu X^M] \end{aligned} \quad (\text{A.6})$$

where ε is an arbitrary field. This would mean that the action has an additional gauge symmetry unless (A.5) holds (for (A.5) implies that the transformations δ_ε are on-shell trivial and thus do not establish additional gauge symmetries). Such gauge symmetries appear to be spurious too. This is suggested by the following toy model which has only one abelian gauge field A_μ and besides is defined through

$$G_{MN} = \left(\begin{array}{cc|c} 0 & 1 & 0 \\ 1 & 0 & 0 \\ \hline 0 & & 1 \end{array} \right), \quad B_{MN} = 0, \quad D = X^1 \quad (M, N = 0, 1, \dots).$$

It is easy to see that, for this model, the general solution to equations (A.2)–(A.4) is, modulo solutions of the form (A.5),

$$\eta^M = \delta_0^M \lambda(X), \quad a_M = -2 \partial_M \lambda(X), \quad b_M = 0$$

where $\lambda(X)$ is a completely arbitrary target space function. This reflects that the action is gauge invariant under

$$\delta_{\bar{\varepsilon}} \gamma_{\mu\nu} = 0, \quad \delta_{\bar{\varepsilon}} X^M = \bar{\varepsilon} \delta_0^M, \quad \delta_{\bar{\varepsilon}} A_\mu = \sqrt{\gamma} \epsilon_{\mu\nu} \partial^\nu \bar{\varepsilon}.$$

[From this one recovers (A.6) for $\bar{\varepsilon} = \varepsilon\lambda$.] As $\delta_{\bar{\varepsilon}}$ generates the shift symmetry $X^0 \rightarrow X^0 + \bar{\varepsilon}$, it is not surprising that the Lagrangian can be cast in a form such that X^0 drops out. Indeed, up to a total derivative the Lagrangian reads

$$L_0 = \sum_{M>1} \frac{1}{2} \sqrt{\gamma} \partial_\mu X^M \partial^\mu X^M - \epsilon^{\mu\nu} \partial_\mu X^1 \cdot (A_\nu - \sqrt{\gamma} \epsilon_{\nu\rho} \partial^\rho X^0),$$

which shows that X^0 can be removed through the field redefinition

$$A'_\mu = A_\mu - \sqrt{\gamma} \epsilon_{\mu\nu} \partial^\nu X^0.$$

Hence, the gauge symmetry (A.6) itself can in this case be removed through a mere local field redefinition and is thus indeed spurious.

B Second part of the cohomological analysis

In this appendix we derive the results described in section 5 by applying $\tilde{s}$ to (3.24) and analysing $\tilde{s}\omega = 0$ modulo $\omega \rightarrow \omega - \tilde{s}\hat{\omega}$ where $\hat{\omega}$ has the same form as ω . We will use the notation

$$\omega = \mathcal{F}(z) + y_\mu^A \mathcal{G}_A^\mu(z) + y_{+-}^\Delta \mathcal{H}_\Delta(z) + y_+^A y_-^B \mathcal{K}_{AB}(z) \quad (\text{B.1})$$

$$y_\mu^A \mathcal{G}_A^\mu(z) = \tilde{X}_+^M \mathcal{G}_M^+(z) + \tilde{Y}_+ \mathcal{G}^+(z) + \tilde{A}_{a+}^{-*} \mathcal{G}^{+a}(z) + (+ \leftrightarrow -) \quad (\text{B.2})$$

$$\begin{aligned} y_{+-}^\Delta \mathcal{H}_\Delta(z) &= \tilde{C}_{a+-}^* \mathcal{H}^a(z) + \tilde{A}_{a+-}^{+*} \mathcal{H}_+^a(z) + \tilde{A}_{a+-}^{-*} \mathcal{H}_-^a(z) \\ &\quad + \tilde{X}_{M+-}^* \mathcal{H}^M(z) + \tilde{X}_{+-}^M \mathcal{H}_M(z) + \tilde{F}_{+-}^a \mathcal{H}_a(z) \end{aligned} \quad (\text{B.3})$$

$$\begin{aligned} y_+^A y_-^B \mathcal{K}_{AB}(z) &= \tilde{X}_+^M \tilde{X}_-^N \mathcal{K}_{MN}(z) + \tilde{X}_+^M \tilde{Y}_- \mathcal{K}_{M-}(z) + \tilde{Y}_+ \tilde{X}_-^M \mathcal{K}_{+M}(z) \\ &\quad + \tilde{X}_+^M \tilde{A}_{a-}^{+*} \mathcal{K}_M^a(z) + \tilde{A}_{a+}^{-*} \tilde{X}_-^M \mathcal{K}_M^a(z) + \tilde{Y}_+ \tilde{Y}_- \mathcal{K}(z) \\ &\quad + \tilde{Y}_+ \tilde{A}_{a-}^{+*} \mathcal{K}_+^a(z) + \tilde{A}_{a+}^{-*} \tilde{Y}_- \mathcal{K}_-^a(z) + \tilde{A}_{a+}^{-*} \tilde{A}_{b-}^{+*} \mathcal{K}^{ab}(z). \end{aligned} \quad (\text{B.4})$$

$\hat{\omega}$ is denoted analogously, with a hat-symbol on the functions of the z 's contained in it.

Note that the order and positions of the superscripts and subscripts is important. In particular, in (B.3) the functions $\mathcal{H}^a$ and $\mathcal{H}_a$ are not related at all, nor are the functions $\mathcal{H}^M$ and $\mathcal{H}_M$ (i.e. they are not related by raising or lowering indices). Furthermore $\mathcal{K}_M^a$ and $\mathcal{K}^a_M$ must not be confused. In the following we will mostly leave out the argument (z) .

Using table 3 in section 5, one obtains

$$\begin{aligned} \tilde{s}\omega &= \left[(\tilde{X}_+^M + \tilde{X}_-^M) \partial_M + \tilde{Y}_\mu \frac{\partial}{\partial \tilde{\Theta}_\mu} + \tilde{F}_{+-}^a \frac{\partial}{\partial \tilde{C}^a} \right] \mathcal{F} \\ &\quad + \tilde{X}_{+-}^M (\mathcal{G}_M^- - \mathcal{G}_M^+) + (\tilde{A}_{a+-}^{+*} - \tilde{X}_-^M \partial_M D_a \cdot) \mathcal{G}^{-a} - (\tilde{A}_{a+-}^{-*} - \tilde{X}_+^M \partial_M D_a \cdot) \mathcal{G}^{+a} \\ &\quad + (-)^{\varepsilon_A} y_+^A \left(\tilde{X}_-^M \partial_M + \tilde{Y}_- \frac{\partial}{\partial \tilde{\Theta}_-} \right) \mathcal{G}_A^+ + (-)^{\varepsilon_A} y_-^A \left(\tilde{X}_+^M \partial_M + \tilde{Y}_+ \frac{\partial}{\partial \tilde{\Theta}_+} \right) \mathcal{G}_A^- \end{aligned}$$

$$\begin{aligned}
& -(\tilde{A}_{a+-}^{+*} + \tilde{A}_{a+-}^{-*})\mathcal{H}^a + (\tilde{X}_+^M \tilde{X}_-^N \partial_M \partial_N D_a + \tilde{X}_{+-}^M \partial_M D_a) \cdot (\mathcal{H}_+^a - \mathcal{H}_-^a) \\
& + \left[-2G_{MN} \tilde{X}_{+-}^N + (H_{KLM} - 2\Gamma_{KLM}) \tilde{X}_+^K \tilde{X}_-^L + \tilde{F}_{+-}^a \partial_M D_a \right] \mathcal{H}^M \\
& + \left[\tilde{X}_+^M y_-^A \mathcal{K}_A^a - (-)^{\varepsilon_A} y_+^A \tilde{X}_-^M \mathcal{K}_A^a \right] \partial_M D_a
\end{aligned} \tag{B.5}$$

where ε_A indicates the Grassmann parity of y_μ^A . $\tilde{s}\omega = 0$ imposes the following conditions:

$$0 = \mathcal{H}^a - \mathcal{G}^{-a} = \mathcal{H}^a + \mathcal{G}^{+a} \tag{B.6}$$

$$0 = \mathcal{H}^M \partial_M D_a + \frac{\partial \mathcal{F}}{\partial \tilde{C}^a} \tag{B.7}$$

$$0 = (\mathcal{H}_+^a - \mathcal{H}_-^a) \partial_M D_a - 2G_{MN} \mathcal{H}^N + \mathcal{G}_M^- - \mathcal{G}_M^+ \tag{B.8}$$

$$0 = \partial_M \mathcal{F} + \mathcal{G}^{+a} \partial_M D_a = \partial_M \mathcal{F} - \mathcal{G}^{-a} \partial_M D_a \tag{B.9}$$

$$0 = \frac{\partial \mathcal{F}}{\partial \tilde{\Theta}_+} = \frac{\partial \mathcal{F}}{\partial \tilde{\Theta}_-} \tag{B.10}$$

$$\begin{aligned}
0 = & (\mathcal{H}_+^a - \mathcal{H}_-^a) \partial_M \partial_N D_a + (H_{MNK} - 2\Gamma_{MNK}) \mathcal{H}^K \\
& - \partial_N \mathcal{G}_M^+ + \partial_M \mathcal{G}_N^- + \mathcal{K}_N^a \partial_M D_a + \mathcal{K}_M^a \partial_N D_a
\end{aligned} \tag{B.11}$$

$$0 = -\frac{\partial \mathcal{G}_M^+}{\partial \tilde{\Theta}_-} + \partial_M \mathcal{G}^- + \mathcal{K}_-^a \partial_M D_a \tag{B.12}$$

$$0 = -\frac{\partial \mathcal{G}_M^-}{\partial \tilde{\Theta}_+} + \partial_M \mathcal{G}^+ - \mathcal{K}_+^a \partial_M D_a \tag{B.13}$$

$$0 = \partial_M \mathcal{G}^{-a} + \mathcal{K}^{ba} \partial_M D_b = \partial_M \mathcal{G}^{+a} - \mathcal{K}^{ab} \partial_M D_b \tag{B.14}$$

$$0 = \frac{\partial \mathcal{G}^+}{\partial \tilde{\Theta}_-} + \frac{\partial \mathcal{G}^-}{\partial \tilde{\Theta}_+} \tag{B.15}$$

$$0 = \frac{\partial \mathcal{G}^{-a}}{\partial \tilde{\Theta}_+} = \frac{\partial \mathcal{G}^{+a}}{\partial \tilde{\Theta}_-} . \tag{B.16}$$

(B.6), (B.9), (B.14) and (B.16) are equivalent to:

$$0 = \partial_M \mathcal{F} - \mathcal{H}^a \partial_M D_a \tag{B.17}$$

$$0 = \partial_M \mathcal{H}^a + \mathcal{K}^{(ab)} \partial_M D_b \tag{B.18}$$

$$0 = \mathcal{K}^{[ab]} \partial_M D_b = 0 \Leftrightarrow \mathcal{K}^{[ab]} = 0 \tag{B.19}$$

$$0 = \frac{\partial \mathcal{H}^a}{\partial \tilde{\Theta}_+} = \frac{\partial \mathcal{H}^a}{\partial \tilde{\Theta}_-} \tag{B.20}$$

where in (B.19) we used (A.1). (B.8) determines $\mathcal{G}_M^+ - \mathcal{G}_M^-$, but not $\mathcal{G}_M^+ + \mathcal{G}_M^- \equiv \mathcal{P}_M$. Hence, (B.8) can be replaced by

$$\mathcal{G}_M^\pm = \frac{1}{2} \mathcal{P}_M \mp G_{MN} \mathcal{H}^N \pm \frac{1}{2} (\mathcal{H}_+^a - \mathcal{H}_-^a) \partial_M D_a . \tag{B.21}$$

Using these expressions, the symmetric and antisymmetric parts of (B.11) read

$$\partial_{(M}[G_{N)K}\mathcal{H}^K] - \Gamma_{MNK}\mathcal{H}^K + \frac{1}{2}\mathcal{A}_{(M}^a\partial_{N)}D_a = 0 \quad (\text{B.22})$$

$$\partial_{[M}\mathcal{P}_{N]} + H_{MNK}\mathcal{H}^K + \mathcal{B}_{[M}^a\partial_{N]}D_a = 0 \quad (\text{B.23})$$

where

$$\begin{aligned} \mathcal{A}_M^a &= \mathcal{K}_M^a + \mathcal{K}_M^a - \partial_M(\mathcal{H}_+^a - \mathcal{H}_-^a) \\ \mathcal{B}_M^a &= \mathcal{K}_M^a - \mathcal{K}_M^a. \end{aligned} \quad (\text{B.24})$$

Using (B.21), Eqs. (B.12) and (B.13) become

$$\begin{aligned} \frac{\partial}{\partial\tilde{\Theta}_-} \left[-G_{MN}\mathcal{H}^N + \frac{1}{2}\mathcal{P}_M + \frac{1}{2}(\mathcal{H}_+^a - \mathcal{H}_-^a)\partial_M D_a \right] &= \partial_M \mathcal{G}^- + \mathcal{K}_-^a \partial_M D_a \\ \frac{\partial}{\partial\tilde{\Theta}_+} \left[G_{MN}\mathcal{H}^N + \frac{1}{2}\mathcal{P}_M - \frac{1}{2}(\mathcal{H}_+^a - \mathcal{H}_-^a)\partial_M D_a \right] &= \partial_M \mathcal{G}^+ - \mathcal{K}_+^a \partial_M D_a. \end{aligned} \quad (\text{B.25})$$

We will now further work out the above equations and remove simultaneously coboundary terms from ω . To this end we examine how the functions of the z 's in ω change under $\omega \rightarrow \omega - \tilde{s}\hat{\omega}$. One finds

$$\mathcal{F} \rightarrow \mathcal{F} \quad (\text{B.26})$$

$$\mathcal{P}_M \rightarrow \mathcal{P}_M - 2\partial_M \hat{\mathcal{F}} + (\hat{\mathcal{G}}^{-a} - \hat{\mathcal{G}}^{+a})\partial_M D_a \quad (\text{B.27})$$

$$\mathcal{G}^\pm \rightarrow \mathcal{G}^\pm - \frac{\partial \hat{\mathcal{F}}}{\partial \tilde{\Theta}_\pm} \quad (\text{B.28})$$

$$\mathcal{H}^a \rightarrow \mathcal{H}^a \quad (\text{B.29})$$

$$\mathcal{H}_\pm^a \rightarrow \mathcal{H}_\pm^a \mp \hat{\mathcal{G}}^{\mp a} + \hat{\mathcal{H}}^a \quad (\text{B.30})$$

$$\mathcal{H}^M \rightarrow \mathcal{H}^M \quad (\text{B.31})$$

$$\mathcal{H}_M \rightarrow \mathcal{H}_M + \hat{\mathcal{G}}_M^+ - \hat{\mathcal{G}}_M^- + 2G_{MN}\mathcal{H}^N - (\hat{\mathcal{H}}_+^a - \hat{\mathcal{H}}_-^a)\partial_M D_a \quad (\text{B.32})$$

$$\mathcal{H}_a \rightarrow \mathcal{H}_a - \hat{\mathcal{H}}^M \partial_M D_a - \frac{\partial \hat{\mathcal{F}}}{\partial \tilde{C}^a} \quad (\text{B.33})$$

$$\begin{aligned} \mathcal{K}_{MN} &\rightarrow \mathcal{K}_{MN} + \partial_N \hat{\mathcal{G}}_M^+ - \partial_M \hat{\mathcal{G}}_N^- - (H_{MKN} - 2\Gamma_{MKN})\hat{\mathcal{H}}^K \\ &\quad - \hat{\mathcal{K}}_N^a \partial_M D_a - \hat{\mathcal{K}}_M^a \partial_N D_a - (\hat{\mathcal{H}}_+^a - \hat{\mathcal{H}}_-^a)\partial_M \partial_N D_a \end{aligned} \quad (\text{B.34})$$

$$\mathcal{K}_{M-} \rightarrow \mathcal{K}_{M-} + \frac{\partial \hat{\mathcal{G}}_M^+}{\partial \tilde{\Theta}_-} - \partial_M \hat{\mathcal{G}}^- - \hat{\mathcal{K}}_-^a \partial_M D_a \quad (\text{B.35})$$

$$\mathcal{K}_{+M} \rightarrow \mathcal{K}_{+M} + \frac{\partial \hat{\mathcal{G}}_M^-}{\partial \tilde{\Theta}_+} - \partial_M \hat{\mathcal{G}}^+ + \hat{\mathcal{K}}_+^a \partial_M D_a \quad (\text{B.36})$$

$$\mathcal{A}_M^a \rightarrow \mathcal{A}_M^a + 2\hat{\mathcal{K}}^{[ab]}\partial_M D_b \quad (\text{B.37})$$

$$\mathcal{B}_M^a \rightarrow \mathcal{B}_M^a - 2\hat{\mathcal{K}}^{(ab)}\partial_M D_b - \partial_M(\hat{\mathcal{G}}^{-a} - \hat{\mathcal{G}}^{+a}) \quad (\text{B.38})$$

$$\mathcal{K} \rightarrow \mathcal{K} - \frac{\partial \hat{\mathcal{G}}^+}{\partial \tilde{\Theta}_-} - \frac{\partial \hat{\mathcal{G}}^-}{\partial \tilde{\Theta}_+} \quad (\text{B.39})$$

$$\mathcal{K}_+^a \rightarrow \mathcal{K}_+^a - \frac{\partial \hat{\mathcal{G}}^{-a}}{\partial \tilde{\Theta}_+} \quad (\text{B.40})$$

$$\mathcal{K}_-^a \rightarrow \mathcal{K}_-^a - \frac{\partial \hat{\mathcal{G}}^{+a}}{\partial \tilde{\Theta}_-} \quad (\text{B.41})$$

$$\mathcal{K}^{(ab)} \rightarrow \mathcal{K}^{(ab)} \quad (\text{B.42})$$

where we have used already (B.21) and (B.24), i.e. we have switched from the functions $\mathcal{G}_M^+ + \mathcal{G}_M^-$, $\mathcal{K}_M^a$ and $\mathcal{K}_M^a$ to $\mathcal{P}_M$, $\mathcal{A}_M^a$, $\mathcal{B}_M^a$. (B.30) and (B.32) show that we can always remove $\mathcal{H}_+^a$, $\mathcal{H}_-^a$ and $\mathcal{H}_M$ completely from ω by choosing $\hat{\mathcal{G}}^{\mp a} = \pm(\mathcal{H}_\pm^a + \hat{\mathcal{H}}^a)$ and $\hat{\mathcal{G}}_M^\mp = \frac{1}{2}\hat{\mathcal{P}}_M \pm \frac{1}{2}\mathcal{H}_M \pm G_{MN}\hat{\mathcal{H}}^N \mp \frac{1}{2}(\hat{\mathcal{H}}_+^a - \hat{\mathcal{H}}_-^a)\partial_M D_a$ where $\hat{\mathcal{P}}_M$ is still arbitrary. In the following we will assume that $\mathcal{H}_+^a$, $\mathcal{H}_-^a$ and $\mathcal{H}_M$ have been removed in this way. Note that this choice of $\hat{\mathcal{G}}^{\mp a}$ and $\hat{\mathcal{G}}_M^\mp$ shifts also $\mathcal{P}_M$, $\mathcal{K}_{MN}$, $\mathcal{K}_{M-}$, $\mathcal{K}_{+M}$, $\mathcal{B}_M^a$, $\mathcal{K}_+^a$ and $\mathcal{K}_-^a$. For simplicity we denote the shifted functions by the same symbols as the original ones. Then we can still work with the above formulae, setting $\mathcal{H}_+^a$, $\mathcal{H}_-^a$ and $\mathcal{H}_M$ to zero everywhere. Hence, without loss of generality we can assume

$$\mathcal{H}_\pm^a = \mathcal{H}_M = 0 \quad (\text{B.43})$$

$$\hat{\mathcal{G}}^{\mp a} = \pm \hat{\mathcal{H}}^a \quad (\text{B.44})$$

$$\hat{\mathcal{G}}_M^\mp = \frac{1}{2}\hat{\mathcal{P}}_M \pm G_{MN}\hat{\mathcal{H}}^N \mp \frac{1}{2}(\hat{\mathcal{H}}_+^a - \hat{\mathcal{H}}_-^a)\partial_M D_a \quad (\text{B.45})$$

where (B.44) and (B.45) guarantee that (B.43) is preserved by the remaining redefinitions (B.26)–(B.42).

In addition to (B.43), we can remove certain parts of other functions occurring in ω through the above redefinitions. To this end we use the superfield notation as in (5.3), leaving out the arguments $(X, \tilde{C})$ of the component functions.

(B.10) and (B.20) require evidently

$$f^\pm = \bar{f} = h^{a\pm} = \bar{h}^a = 0 \quad \Leftrightarrow \quad \mathcal{F} = f, \quad \mathcal{H}^a = h^a \quad (\text{B.46})$$

(B.15) requires $g^{+-} = -g^{-+}$ and $\bar{g}^+ = \bar{g}^- = 0$. Furthermore, (B.28) gives $g^\pm \rightarrow g^\pm - \hat{f}^\pm$ and $g^{+-} \rightarrow g^{+-} - \bar{\hat{f}}$. Hence, we can always remove g^+ , g^- , g^{+-} and g^{-+} from ω by choosing $\hat{f}^\pm$ and $\bar{\hat{f}}$ appropriately. By the same reasoning that has led to (B.43)–(B.45) we can therefore assume, without loss of generality,

$$g^\pm = g^{+-} = g^{-+} = \bar{g}^\pm = 0 \quad \Leftrightarrow \quad \mathcal{G}^\pm = \tilde{\Theta}_\pm g^{\pm\pm} \quad (\text{B.47})$$

$$\hat{f}^\pm = \bar{\hat{f}} = 0 \quad \Leftrightarrow \quad \hat{\mathcal{F}} = \hat{f} \quad (\text{B.48})$$

Analogously we conclude from (B.39)–(B.41), taking (B.44) into account, that without loss of generality we can assume

$$k = k^\pm = 0 \quad \Leftrightarrow \quad \mathcal{K} = \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k} \quad (\text{B.49})$$

$$\bar{\hat{g}}^\pm = 0, \quad \hat{g}^{+-} = -\hat{g}^{-+} \equiv \hat{g} \quad \Leftrightarrow \quad \hat{\mathcal{G}}^\pm = \hat{g}^\pm \pm \tilde{\Theta}_\mp \hat{g} + \tilde{\Theta}_\pm \hat{g}^{\pm\pm} \quad (\text{B.50})$$

and

$$\begin{aligned}
k_+^a = k_-^a = \hat{h}^{a\pm} = \bar{\hat{h}}^a = 0, \quad k_-^{a+} = -k_+^{a-} \equiv k^a \\
\Leftrightarrow \begin{cases} \mathcal{K}_+^a = -\tilde{\Theta}_- k^a + \tilde{\Theta}_+ k_+^{a+} + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_+^a \\ \mathcal{K}_-^a = \tilde{\Theta}_+ k^a + \tilde{\Theta}_- k_-^{a-} + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_-^a \\ \hat{\mathcal{H}}^a = \hat{h}^a \end{cases} \quad (B.51)
\end{aligned}$$

Taking into account (B.45) and (B.50), Eqs. (B.35) and (B.36) yield analogously

$$\begin{aligned}
k_{M-} = k_{+M} = 0, \quad k_{M-}^+ = k_{+M}^- \equiv k_M \\
\Leftrightarrow \begin{cases} \mathcal{K}_{M-} = \tilde{\Theta}_+ k_M + \tilde{\Theta}_- k_{M-}^- + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{M-} \\ \mathcal{K}_{+M} = \tilde{\Theta}_- k_M + \tilde{\Theta}_+ k_{+M}^+ + \tilde{\Theta}_+ \tilde{\Theta}_- \bar{k}_{+M} \end{cases}, \quad (B.52)
\end{aligned}$$

as well as Eqs. (5.13)–(5.15) up to an extra contribution $2\partial_M \hat{g}$ to $\bar{\hat{p}}_M$ which can be neglected in (5.15) as it provides an $\tilde{s}$ -exact contribution to $\hat{\omega}$ (furthermore we have used the notation $\hat{\mathcal{H}}_+^a - \hat{\mathcal{H}}_-^a = 2\hat{Z}^a$ in section 5).

Next we analyse Eqs. (B.25), using (B.43), (B.47) and (B.51). The $\tilde{\Theta}$ independent components of Eqs. (B.25) yield

$$p_M^\pm = \mp 2G_{MN} h^{N\pm}, \quad (B.53)$$

the terms linear in the $\tilde{\Theta}$'s give

$$\bar{p}_M = 0 \quad (B.54)$$

$$G_{MN} \bar{h}^N = k^a \partial_M D_a \quad (B.55)$$

$$\partial_M g^{\pm\pm} = \pm k_\pm^{a\pm} \partial_M D_a, \quad (B.56)$$

and the $\tilde{\Theta}_+ \tilde{\Theta}_-$ components yield

$$\bar{k}_-^a \partial_M D_a = \bar{k}_+^a \partial_M D_a = 0 \quad \Leftrightarrow \quad \bar{k}_-^a = \bar{k}_+^a = 0 \quad (B.57)$$

where we used (A.1).

Inserting (B.46) in (B.18) we get in particular

$$k^{(ab)\pm} \partial_M D_a = \bar{k}^{(ab)} \partial_M D_a = 0 \quad \Leftrightarrow \quad k^{(ab)\pm} = \bar{k}^{(ab)} = 0 \quad (B.58)$$

where we used again (A.1).

Finally we examine the $\tilde{\Theta}_+ \tilde{\Theta}_-$ components of (B.22), (B.23) and (B.7). Using (B.46), (B.54) and (B.55) one gets

$$\begin{aligned}
\partial_{[M} [k^a \partial_{N]} D_a] - \Gamma_{MNK} \bar{h}^K + \frac{1}{2} \bar{a}_{(M}^a \partial_{N]} D_a &= 0 \\
H_{MNK} \bar{h}^K + \bar{b}_{[M}^a \partial_{N]} D_a &= 0 \\
\bar{h}^M \partial_M D_a = 0, \quad G_{MN} \bar{h}^N = k^a \partial_M D_a. & \quad (B.59)
\end{aligned}$$

Due to (A.2)–(A.5) we conclude from (B.59), using (B.37) and (B.38), that we can assume without loss of generality

$$\bar{h}^M = k^a = \bar{a}_M^a = \bar{b}_M^a = \bar{\bar{k}}^{ab} = 0. \quad (B.60)$$

Altogether this provides the results given in section 5.

C Nonabelian deformations

In this appendix we describe some obvious deformations arising from (6.21) and show how nonabelian Born-Infeld type actions emerge from them in special cases.

Assume we choose a Lagrangian (4.9) which is invariant under the global action of a compact Lie group represented linearly on the X^M and A_μ^a , such that the representation of the A_μ^a is the adjoint one (i.e., the indices a of the abelian gauge fields label simultaneously the generators of the Lie group). That is to say, assume

$$\delta_a X^M = -T_{aN}^M X^N, \quad \delta_a A_\mu^b = -f_{ac}^b A_\mu^c \quad \Rightarrow \quad \delta_a L = 0$$

where δ_a denote the generators of the Lie group, f_{bc}^a are the structure constants of the corresponding Lie algebra, and $\{T_a\}$ is the matrix representation of that Lie algebra on the X^M ,

$$[T_a, T_b] = f_{ab}^c T_c.$$

This requires of course the functions $G_{MN}(X)$, $B_{MN}(X)$ and $D_a(X)$ to transform under $\mathcal{G}$ according to the representation indicated by their indices. Then it is clear that a nonabelian deformation of the Lagrangian is obtained just by the replacements

$$\begin{aligned} \partial_\mu A_\nu^a - \partial_\nu A_\mu^a &\rightarrow \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f_{bc}^a A_\mu^b A_\nu^c \\ \partial_\mu X^M &\rightarrow D_\mu X^M = \partial_\mu X^M + g A_\mu^a T_{aN}^M X^N \end{aligned}$$

where g is a constant deformation parameter. The deformed Lagrangian is invariant under deformed gauge transformations

$$\delta_\varepsilon X^M = -g \varepsilon^a T_{aN}^M X^N, \quad \delta_\varepsilon A_\mu^a = \partial_\mu \varepsilon^a + g f_{bc}^a \varepsilon^b A_\mu^c.$$

To first order in g , these deformations are indeed given by (6.21) with

$$h_{bc}^a = f_{bc}^a, \quad h_a^M = -T_{aN}^M X^N, \quad p'_{Ma} = b_{bM}^a = a_{bM}^a = k_{cd}^{(ab)} = 0.$$

This holds as $\mathcal{L}_a G_{MN} = \mathcal{L}_a B_{MN} = 0$ are implied by the assumption that $G_{MN}(X)$ and $B_{MN}(X)$ transform under $\mathcal{G}$ as indicated by their indices.

Nonabelian Born-Infeld type actions are obtained from the above formulae as follows. We split the X^M into two subsets,

$$\{X^M\} = \{x^m, \varphi^a\},$$

where the φ^a transform according to the adjoint representation of some compact Lie group. For simplicity we assume that the x^m transform according to the trivial representation, i.e.

$$T_{ab}^c = -f_{ab}^c, \quad T_{am}^n = 0.$$

We now choose a faithful matrix representation $\{\mathbf{t}_a\}$ of the Lie algebra and introduce the matrix

$$\varphi = \varphi^a \mathbf{t}_a.$$

Next we construct a Lagrangian (4.9) with

$$G_{mn}(X) = g_{mn}(x) \text{Tr}(\mathbf{1} - \boldsymbol{\varphi}^2)^{-1/2}, \quad G_{ma} = G_{ab} = B_{MN} = 0$$

$$D_a(X) = -\text{Tr}[\mathbf{t}_a \boldsymbol{\varphi} (\mathbf{1} - \boldsymbol{\varphi}^2)^{-1/2}]$$

where $(\mathbf{1} - \boldsymbol{\varphi}^2)^{-1/2}$ is understood as a power series in $\boldsymbol{\varphi}$. A nonabelian deformation of the corresponding Lagrangian is now given by

$$L_0^{\text{nonabel.}} = \frac{1}{2} \text{Tr} \left\{ (\mathbf{1} - \boldsymbol{\varphi}^2)^{-1/2} (\mathbf{1} \sqrt{\gamma} \gamma^{\mu\nu} g_{mn}(x) \partial_\mu x^m \cdot \partial_\nu x^n - \epsilon^{\mu\nu} \boldsymbol{\varphi} \mathbf{F}_{\mu\nu}) \right\}$$

where

$$\mathbf{F}_{\mu\nu} = (\partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f_{bc}^a A_\mu^b A_\nu^c) \mathbf{t}_a .$$

Elimination of the auxiliary fields $\gamma_{\mu\nu}$ and φ^a yields (6.33). Note that this example can be easily generalized by allowing the x^m to transform linearly under the Lie group, or nontrivial B_{mn} in the way outlined above. Note also that an ordering ambiguity of matrices under the above traces does not arise.

References

- [1] See J. Polchinski, Tasi Lectures on D-branes, hep-th/9611050 and refs. therein.
- [2] P. K. Townsend, *Phys. Lett.* **409B** (1997) 131.
- [3] M. Cederwall and P. K. Townsend, JHEP 09 (1997) 003 (hep-th/9709002);
M. Cederwall and A. Westerberg, World volume fields, SL(2,Z) and duality: the type IIB three-brane, hep-th/9710007.
- [4] J. Gomis and S. Weinberg, *Nucl. Phys.* **B469** (1996) 473.
- [5] G. Barnich, F. Brandt and M. Henneaux, *Commun. Math. Phys.* **174** (1995) 57.
- [6] G. Barnich and M. Henneaux, *Phys. Lett.* **311B** (1993) 123.
- [7] This was first observed in the pioneering works C. Becchi, A. Rouet and R. Stora, *Phys. Lett.* **52B** (1974) 344; *Commun. Math. Phys.* **42** (1975) 127; *Ann. Phys.* **98** (1976) 287.
See also [9, 10] and refs. therein.
- [8] I. A. Batalin and G. A. Vilkovisky, *Phys. Lett.* **102B** (1981) 27; *Phys. Rev.* **D28** (1983) 2567 (E: **D30** (1984) 508).
- [9] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton, 1992).
- [10] J. Gomis, J. París and S. Samuel, *Phys. Rep.* **259** (1995) 1.

- [11] J. M. L. Fisch and M. Henneaux, *Commun. Math. Phys.* **128** (1990) 627;
M. Henneaux, *Nucl. Phys. B (Proc. Suppl.)* **18A** (1990) 47.
See also [9].
- [12] F. Brandt, *Commun. Math. Phys.* **190** (1997) 459.
- [13] F. Brandt, W. Troost and A. Van Proeyen, *Nucl. Phys.* **B464** (1996) 353.
- [14] F. Brandt, J. Gomis and J. Simón, *Phys. Lett.* **419B** (1998) 148.
- [15] F. Brandt, M. Henneaux and A. Wilch, *Nucl. Phys. (PM)* **B510** (1998) 640.
- [16] B. de Wit and P. van Nieuwenhuizen, *Nucl. Phys.* **B312** (1989) 58;
C. M. Hull and B. Spence, *Phys. Lett.* **232B** (1989) 204.
- [17] A. A. Tseytlin, *Nucl. Phys.* **B469** (1996) 51.
- [18] F. Brandt, W. Troost and A. Van Proeyen, *Phys. Lett.* **374B** (1996) 31.
- [19] F. Brandt, N. Dragon and M. Kreuzer, *Phys. Lett.* **231B** (1989) 263; *Nucl. Phys.*
B332 (1990) 224; *Nucl. Phys.* **B340** (1990) 187.
- [20] F. Brandt, *Phys. Lett.* **320B** (1994) 57;
G. Barnich and M. Henneaux, *Phys. Rev. Lett.* **72** (1994) 1588;
G. Barnich, F. Brandt and M. Henneaux, *Commun. Math. Phys.* **174** (1995) 93;
Nucl. Phys. **B455** (1995) 357.